



This is a repository copy of *Perioperative laboratory monitoring in congenital haemophilia patients with inhibitors*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/150119/>

Version: Accepted Version

Article:

Hart, D.P., Hay, C.R.M., Liesner, R. et al. (3 more authors) (2019) Perioperative laboratory monitoring in congenital haemophilia patients with inhibitors. *Blood Coagulation & Fibrinolysis*, 30 (7). pp. 309-323. ISSN 0957-5235

<https://doi.org/10.1097/mbc.0000000000000840>

Article available under the terms of the CC-BY-NC licence
(<https://creativecommons.org/licenses/by-nc/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Blood Coagulation & Fibrinolysis

Perioperative Laboratory Monitoring in Congenital Haemophilia Patients with Inhibitors: A Systematic Literature Review --Manuscript Draft--

Manuscript Number:	BCF-19-064R1
Full Title:	Perioperative Laboratory Monitoring in Congenital Haemophilia Patients with Inhibitors: A Systematic Literature Review
Article Type:	Review Article/Proposal
Section/Category:	Clinical study
Additional Information:	
Question	Response
Keywords:	Haemophilia; Surgery; Inhibitors; Laboratory Monitoring; Systematic Literature Review
Manuscript Classifications:	2.2: Haemophilias (including inhibitors) and other rare factor disorders; 2.3: Inhibitors; 3.2: Perioperative bleeding; 9.2: Routine clinical
Corresponding Author:	Maria Haughton, MSci Costello Medical Consulting Ltd Cambridge, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Costello Medical Consulting Ltd
Corresponding Author's Secondary Institution:	
First Author:	Daniel P Hart
First Author Secondary Information:	
Order of Authors:	Daniel P Hart
	Charles RM Hay
	Ri Liesner
	Guillermo Tobaruela
	Bethan Du-Mont
	Mike Makris
	Maria Haughton, MSci
Order of Authors Secondary Information:	
Manuscript Region of Origin:	UNITED KINGDOM
Suggested Reviewers:	
Opposed Reviewers:	
Abstract:	Although the use of clotting factor concentrates is the mainstay of haemophilia care, the development of inhibitors complicates disease management. Perioperative management of patients with inhibitors is therefore a challenge. A systematic literature review was performed to identify literature reporting on the perioperative monitoring and management of haemophilia. MEDLINE, Embase and Cochrane databases were searched from database inception to 26 March 2018. Recent congress proceedings were also searched. Titles and abstracts, then full texts, were screened for relevance by two reviewers. Quality of included studies was assessed using the Critical Appraisal Skills Programme checklist. Of the 2,033 individual entries identified, 86 articles met the inclusion criteria. The identified studies were screened again to find papers

	<p>reporting perioperative laboratory monitoring in patients with congenital haemophilia A or B, resulting in 24 articles undergoing data extraction. Routine perioperative assay monitoring practices were the most commonly reported (n=20/24); thrombin generation assay (TGA) was the least commonly reported (n=2/24). Other monitoring practices described were factor VII and factor VIII coagulation activity (FVII:C, FVIII:C) (n=8/24, n=5/24, respectively), and thromboelastography (TEG) or rotational thromboelastometry (ROTEM) assessments (n=3/24). The impact of monitoring on treatment decisions was, however, rarely reported. In conclusion, many methods of perioperative monitoring of haemophilia patients with inhibitors have been identified in this review, yet there is a lack of reporting in larger scale cohort studies. More detailed reporting on the impact of monitoring outcomes on treatment decisions is also needed to share best practice, particularly as new therapeutic agents emerge.</p>
--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Perioperative Laboratory Monitoring in Congenital
Haemophilia Patients with Inhibitors: A Systematic
Literature Review**

Daniel P Hart¹, Charles RM Hay², Ri Liesner³, Guillermo Tobaruela⁴, Bethan Du-Mont⁵,
Mike Makris⁶

¹The Royal London Hospital Haemophilia Centre, Barts and The London School of
Medicine and Dentistry, Queen Mary University of London, London, UK; ²Central
Manchester University Hospitals Foundation Trust, Manchester, UK; ³Great Ormond
Street Hospital for Children NHS Foundation Trust, London, UK; ⁴Roche Products
Ltd., Welwyn Garden City, UK; ⁵Costello Medical, Cambridge, UK; ⁶Sheffield
Haemophilia and Thrombosis Centre, Royal Hallamshire Hospital, Sheffield, UK

Correspondence to: Daniel P Hart

Postal Address: The Royal London Hospital Haemophilia Centre, Barts and
The London School of Medicine and Dentistry, Queen Mary University of
London, London, UK
Email Address: d.hart@qmul.ac.uk

Short title: Perioperative Haemophilia Monitoring

Funding: Roche Products Ltd. and Chugai Pharma UK Ltd.

Abstract

Although the use of clotting factor concentrates is the mainstay of haemophilia care, the development of inhibitors complicates disease management. Perioperative management of patients with inhibitors is therefore a challenge. A systematic literature review was performed to identify literature reporting on the perioperative monitoring and management of haemophilia. MEDLINE, Embase and Cochrane databases were searched from database inception to 26 March 2018. Recent congress proceedings were also searched. Titles and abstracts, then full texts, were screened for relevance by two reviewers. Quality of included studies was assessed using the Critical Appraisal Skills Programme checklist. Of the 2,033 individual entries identified, 86 articles met the inclusion criteria. The identified studies were screened again to find papers reporting perioperative laboratory monitoring in patients with congenital haemophilia A or B, resulting in 24 articles undergoing data extraction. Routine perioperative assay monitoring practices were the most commonly reported (n=20/24); thrombin generation assay (TGA) was the least commonly reported (n=2/24). Other monitoring practices described were factor VII and factor VIII coagulation activity (FVII:C, FVIII:C) (n=8/24, n=5/24, respectively), and thromboelastography (TEG) or rotational thromboelastometry (ROTEM) assessments (n=3/24). The impact of monitoring on treatment decisions was, however, rarely reported. In conclusion, many methods of perioperative monitoring of haemophilia patients with inhibitors have been identified in this review, yet there is a lack of reporting in larger scale cohort studies. More detailed reporting on the impact of monitoring outcomes on treatment decisions is also needed to share best practice, particularly as new therapeutic agents emerge.

Key words: Haemophilia; Surgery; Inhibitors; Laboratory Monitoring; Systematic Literature Review

1 INTRODUCTION

2 Haemophilia is a rare disease caused by a deficiency of coagulation factor VIII
3 (FVIII) (haemophilia A, HA) or factor IX (FIX) (haemophilia B, HB) and leaves
4 patients more prone to excessive bleeding.[1] The standard of care for severe
5 haemophilia (factor activity <1 IU/dl), and some patients with moderate haemophilia
6 (factor activity 1-5 IU/dl), is to prevent or minimise bleeding episodes using infusions
7 of the missing factor concentrate (prophylaxis). Breakthrough bleeds may still occur
8 requiring additional on-demand treatment. Treatment protocols, intensity of
9 prophylaxis or choice to remain on-demand alone must be tailored to individual
10 needs together with consideration of the local health economics.[1]

11 In response to regular treatment, a subgroup of haemophilia patients of all severities
12 can produce immunoglobulin G (IgG) antibodies, termed 'inhibitors', which work to
13 neutralise clotting factors.[2, 3] Inhibitors complicate prophylaxis and on-demand
14 management by reducing or fully neutralising the efficacy of infused factor
15 concentrates, depending on the detected inhibitor titre.[1, 4]

16 Persistent inhibitors are a concern for haemophilia patients, particularly if undergoing
17 surgical procedures.[5] Permanent tolerance induction in severe haemophilia
18 (immune tolerance induction, ITI) is the preferable strategy to minimise future
19 bleeding and/or management risks of surgery in the presence of an inhibitor.[6, 7]
20 Tolerising practices may also include the use of anti-CD20 monoclonal antibody,
21 immunoadsorption and plasmapheresis for additional short-term benefit.[5]
22 Strategies for re-achieving tolerance in non-severe HA are less well defined.[8]

23 Knowledge of previous and/or current inhibitor status prior to surgery is crucial,
24 either as repeat laboratory assessment ahead of surgery if time allows (severe
25 haemophilia), reference to laboratory screening since the most recent FVIII
26 concentrate exposure (non-severe HA), or attention to *in vivo* recovery and
27 perioperative efficacy of infused concentrate (all severities) during the peri- and
28 post-surgical course.[6, 9]

29 The monitoring and management of haemophilia patients with inhibitors undergoing
30 surgical procedures is a particular challenge, and is the focus of this review.

31 MATERIALS AND METHODS

32 Search Strategy for Identification of Studies

1 A systematic literature review was performed in accordance with a pre-specified
2 search protocol designed to identify literature reporting on the perioperative
3 monitoring and management of haemophilia patients with inhibitors. The review
4 process involved searching electronic databases, and hand-searching of key
5 haemophilia/haematology conference proceedings from the last two years and
6 reference lists of any relevant systematic reviews identified during the searches.

7 All electronic databases were searched on 26 March 2018. The databases searched
8 to identify relevant published literature were: MEDLINE, MEDLINE In-Process,
9 MEDLINE Daily and MEDLINE Epub Ahead of Print (1946 to present); Embase (1974
10 to 23 March 2018); The Cochrane Database of Systematic Reviews (CDSR): Issue 3
11 of 12, March 2018; The Database of Abstracts of Reviews of Effects (DARE): Issue 2
12 of 4, April 2015; The Cochrane Central Register of Controlled Trials (CENTRAL):
13 Issue 2 of 12, February 2018.

14 In addition to the electronic database searches, hand-searches were performed to
15 generate further evidence from a variety of sources. The bibliographies of published
16 systematic reviews identified through the electronic database searches were hand-
17 searched to identify any additional relevant studies for inclusion in the review. The
18 proceedings of relevant haemophilia and haematology congresses that had taken
19 place within approximately two and a half years prior to December 2017 were also
20 hand-searched, including: American Society of Hematology Annual Meeting (2015,
21 2016, 2017); British Society for Haematology Annual Scientific Meeting (2016, 2017);
22 European Association for Haemophilia and Allied Disorders Annual Congress (2016,
23 2017); Haemophilia and Thrombosis Research Society Scientific Symposium (2015,
24 2017); European Hematology Association (EHA) Congress (2016, 2017);
25 International Society for Thrombosis and Haemostasis Congress (2016, 2017); World
26 Federation of Hemophilia (WFH) World Congress (2016). The websites and abstract
27 books of these congresses were searched, if available, using terms based on the
28 complete list of electronic database search terms.

29 Full details of the search strategies for the electronic database searches and the
30 congress searches are presented in Supplementary Table 1–3.

31 **Study Selection**

32 All articles retrieved through the electronic database searches and hand-searches
33 were screened by two independent reviewers and included based on their alignment

with the predefined eligibility criteria (Supplementary Table 4). The strategy was specifically designed to capture studies reporting on the monitoring and/or management of haemophilia patients with inhibitors undergoing surgery. All articles were initially screened based on their abstracts only. Following the abstract review stage, the full texts of remaining articles were then screened for relevance to produce the final list of included studies. For pragmatic reasons, additional study design eligibility criteria were applied during the screening of full texts to ensure that only interventional and observational studies were included in the final list. Study designs were determined by their description as reported by authors in the papers. In cases where the study design was not explicitly stated, the reviewers defined observational studies as those that examined and analysed the data of a patient cohort as a group, compared to an analysis of individual patient data only in case series. Studies that described patients being assigned to a specific treatment group were categorised as interventional. A full list of papers excluded, and the reasons for their exclusion, at the full text screening stages of the review is shown in Supplementary Table 5. In addition, a list of the case studies identified during the review and excluded based on study design prior to full text screening is available in Supplementary Table 6.

Data Extraction and Analysis

Following application of the eligibility criteria, there was still a large number of included studies. Therefore, the list was screened an additional time by one reviewer, and checked by a second reviewer, to identify the studies that reported perioperative laboratory monitoring in patients with congenital HA or HB. In cases of studies reporting on both congenital and acquired haemophilia patients, only information on congenital patients was extracted. Data were extracted from each article by a single individual, and reviewed by a second.

Since included studies are of different designs, their quality was assessed using the Critical Appraisal Skills Programme (CASP) checklist most appropriate for the study design (e.g. case control study, cohort study, randomised controlled trial).

RESULTS

Search Results

The literature search retrieved 1,481 abstracts from electronic databases, 502 abstracts from conference proceedings and 50 articles from hand searches of

existing review bibliographies. Following application of the eligibility criteria to the identified abstracts and, subsequently, full text articles (Figure 1), a final list of 86 relevant articles was identified (Table 1).

In the interest of focusing the review more specifically, the 86 full text articles were screened once more to identify the studies reporting perioperative laboratory monitoring in patients with congenital HA or HB and inhibitors. This resulted in a final list of 24 articles that underwent full data extraction (Table 2).

Quality Assessment

Quality assessments were carried out for the interventional and observational studies that underwent data extraction (Supplementary Table 7–8). Overall, the issues addressed by the interventional studies were clearly focused and all patients were properly accounted for. The most substantial limitation of the interventional studies was a lack of blinding (8/9 studies), which may have led to bias in the ascertainment and reporting of outcomes. For the observational studies, follow-up was almost always complete (14/15) but due to a lack of reporting, many of the quality assessment questions had to be marked as not applicable.

Perioperative Monitoring

Overall, 40% (34/86) of articles identified through the review mentioned how patients were monitored; of these, nearly three quarters (24) of the articles mentioned the use of laboratory monitoring in patients with inhibitors complicating congenital HA or HB, either in the methods section or when describing the outcomes of the study (Table 1). However, even in studies which mentioned laboratory monitoring, the use and impact of specific monitoring protocols on treatment decisions was often not well described. Instead, it was common for the details of perioperative monitoring to be provided for information only, or to report the haemostatic efficacy of the treatment.

Routine laboratory monitoring

Amongst the studies reporting perioperative laboratory monitoring practices, routine laboratory monitoring practices, such as platelet count, prothrombin time (PT), activated partial thromboplastin time (APTT), fibrinogen levels, D-dimer levels and antithrombin (ATIII) were the most commonly reported. This monitoring information

was frequently provided to demonstrate efficacy, or lack thereof, and determine safety, of the haemostatic agent, particularly in the context of interventional studies.[10, 11] In the majority of studies it was unclear whether the outcomes of laboratory tests were available to the care team within the timeframe necessary to influence treatment decisions (Table 3).[12, 13]

When the influence of monitoring on clinical decisions was discussed, this was mainly in the context of individual patient cases, such as reduction in activated prothrombin complex concentrate (aPCC) treatment following elevation in D-dimer levels,[14] as opposed to providing insight into how laboratory monitoring influenced treatment decisions and outcomes on a cohort-wide level. In other cases, laboratory tests were only utilised in patients who experienced an adverse event.[15]

Overall, very limited information was found in the identified studies to indicate the influence of perioperative monitoring results on clinical decisions.

Factor VII

A total of 8 of the 17 extracted studies describing treatment with recombinant factor VIIa (rFVIIa), administered continually (CI) or using bolus doses, described the monitoring of factor VII coagulation activity (FVII:C) (Table 4). Two studies published by the same centre described monitoring FVIIa levels using a one-stage coagulation assay suggesting that coagulation activity as opposed to protein levels was assessed. The studies from this centre also report that the specific FVIIa assay (Staclot™, Diagnostica Stago) was not found to be practical or reliable.[16, 17]

Even where FVII:C monitoring is mentioned, FVII:C was rarely used to make dosing decisions. In one study investigating continuous infusion of rFVIIa, observed bleeding was used to deduce that the target FVII:C of 10 IU/dl was insufficient.[18] Two other studies noted that in patients with ineffective haemostatic efficacy, FVII:C often exceeded the target 30 IU/ml.[10, 19]

When comparing CI to bolus administration, one study found that FVII:C levels were consistently higher in patients undergoing CI; however, the difference was not statistically significant, and there was no difference in haemostatic efficacy between groups (75% vs. 73%, respectively).[19]

Thromboelastography/Rotational thromboelastography analysis

Thromboelastography (TEG) or rotational thromboelastography (ROTEM) coagulation assessments were used in three of the more recent studies assessing a variety of agents (including rFVIIa, aPCC, and FVIII) (Table 5). In one study, ROTEM analysis on *in vitro* samples was used to identify the minimum necessary dose of activated coagulation products and most suitable treatment (rFVIIa vs. aPCC) for perioperative haemostatic control in patients with inhibitors.[20] This study found that preoperative *in vitro* ROTEM analysis more accurately predicted the impact of treatment with rFVIIa than with aPCC.[20] The other studies used ROTEM intraoperatively to demonstrate haemostatic efficacy, but did not report how the outcomes impacted clinical treatment decisions.[21, 22]

Factor VIII:C

Four studies were found to have analysed human FVIII:C, while one analysed porcine FVIII:C (Table 6).[13, 21-24] The use of FVIII:C to identify cases of 'resistance' to porcine FVIII concentrate was discussed in an early study,[13] however, only one recent study mentioned how FVIII:C monitoring influenced treatment decisions.[24] This retrospective observational study involved routine monitoring of FVIII:C and when one patient's FVIII:C levels declined after a total knee replacement, their treatment was switched from cryoprecipitate to plasma-derived FVIII (pdFVIII), leading to a good haemostatic outcome.[24]

Thrombin generation

Only two articles described monitoring using a thrombin generation assay (TGA) (Table 7).[22, 25] In one of these studies, TGA parameters were used to assess haemostatic efficacy, with no indication of how the results impacted care decisions.[22] The other, a 2016 study involving 10 inhibitor patients undergoing orthopaedic surgery, investigated the association between TGA and clinical bleeding events, finding that there was no difference in the TGA values between patients who did and did not experience bleeding complications.[25]

DISCUSSION

Whilst this review uncovered evidence on the methods of monitoring and management used in haemophilia patients undergoing surgery, there was very little information to indicate how the outcomes of laboratory monitoring practices were

1 used to influence treatment decisions. In papers where the impact of monitoring was
2 mentioned, this tended to be described on an individual patient basis. Therefore, it is
3 difficult to use the available evidence to understand to what extent laboratory
4 monitoring is used in clinical practice and how it could be utilised to improve patient
5 care.

6 Another barrier to understanding the impact of perioperative monitoring in
7 haemophilia patients is the lack of a generalisable assay for aPCC and rFVIIa.
8 Monitoring of patients undergoing treatment with these agents is currently
9 conducted according to local protocols, instead of a global standard, with centres
10 forced to evaluate treatment efficacy through clinical, rather than laboratory,
11 assessments.

12 In addition, while some evidence related to monitoring and management with
13 traditional treatments, such as rFVIIa, aPCC and FVIII was found, the review did not
14 identify published literature reporting on such practices in patients treated with
15 emerging therapeutics, such as emicizumab, concizumab and fitusiran. There are
16 currently only anecdotal data about surgical haemostasis planning in patients with
17 these agents on board.[26, 27] In such scenarios, monitoring for all these agents will
18 be complex, both in terms of consideration of appropriate laboratory assays,
19 together with interpreting results in the context of global haemostatic potential.

20 Emicizumab, a recombinant, humanised, bispecific monoclonal antibody recently
21 approved for treatment of HA with inhibitors by the Food and Drug Administration
22 (FDA), works by acting similarly to activated factor VIII in bridging activated factor
23 IX and factor X to trigger the coagulation cascade.[28] As emicizumab affects
24 intrinsic pathway clotting-based laboratory tests, including activated clotting time
25 (ACT) and all assays based on APTT, intrinsic pathway clotting-based laboratory test
26 results should not be used to monitor emicizumab activity, determine dosing for
27 factor replacement or anti-coagulation, or measure FVIII inhibitor titres in patients
28 receiving this treatment.[29] A recent case study describing the use of rescue aPCC
29 treatment to provide additional haemostatic control during a spontaneous bleeding
30 event in a patient receiving prophylactic emicizumab used TGA to determine the
31 optimal aPCC dosage to maintain haemostatic efficacy while limiting the risk of
32 thrombotic complications.[30] While this review identified an interventional study
33 reporting the use of emicizumab in haemophilia patients with inhibitors undergoing
34 surgery, no information on perioperative laboratory monitoring was provided.[27]

1 Concizumab, a humanized monoclonal antibody against tissue factor pathway
2 inhibitor, is under investigation for the prophylactic treatment of HA and HB patients
3 with inhibitors in a phase II trial due to complete in 2019 (NCT03196284).[31]
4 Previous studies have used TGA to demonstrate concizumab activity, suggesting that
5 this may be a potential method of evaluating efficacy in clinical practice.[32] Lastly,
6 fitusiran, an investigational RNA interference therapy, works by targeting ATIII
7 messenger RNA to suppress ATIII production.[33] One interventional study
8 examining fitusiran treatment in haemophilia patients undergoing surgery was
9 identified in this review, but no details on perioperative laboratory monitoring were
10 reported.[34] As these emerging therapies enter the market it will be important to
11 establish an understanding of the standard of care used to monitor patients receiving
12 these treatments.

13 While data on these emerging therapies are currently limited it is difficult to predict
14 response to treatment during surgery, as well as expectations from monitoring.
15 Efforts have been made to standardise TGA, but these developments are rarely
16 shared outside of research settings. More information is therefore needed to
17 understand their use in patients with inhibitors undergoing surgery in clinical
18 practice. As the availability of data on suitable assays remains limited, the United
19 Kingdom Haemophilia Centres Doctors' Organisation has recommended that non-
20 urgent major surgery in haemophilia patients with inhibitors receiving novel
21 prophylaxis agents is delayed, until more specific methods of monitoring these
22 patients can be found or treatment algorithms and risks are better understood.[35]

23 For pragmatic reasons, the scope of the review was limited to include only
24 interventional and observational studies. The results of this review indicate that the
25 currently available higher quality evidence base provides little insight into the
26 standard of care for the use of laboratory monitoring in the management of
27 haemophilia patients with inhibitors undergoing surgery. This topic has also been
28 addressed in case studies and series, including a case series reported by Dargaud et
29 al. in 2010.[36] This paper investigated the use of TGA in monitoring efficacy of
30 agents in surgical procedures for six patients and showed that TGA results correlated
31 with clinical bleeding risk and endogenous thrombin potential can be used to monitor
32 agent efficacy in their hands.[36] These results were not supported by a later, larger
33 scale study involving 10 inhibitor patients undergoing surgery identified by our
34 review however, as no significant differences in TGA parameters between

haemophilia patients with inhibitors who did and did not experience bleeding complications following surgery were found.[25] Whilst case studies and series can provide valuable insight, they are considered to be a source of lower quality evidence due to their risk of bias and potential lack of generalisability to a wider patient population. The results of this review, therefore, highlight the need to report higher quality evidence on monitoring and managing surgical haemophilia patients with inhibitors to establish a standard of care in this area.

CONCLUSION

In conclusion, this systematic literature review demonstrated that there are multiple methods of laboratory monitoring used in haemophilia patients with inhibitors undergoing surgery, although these are largely reported in the context of clinical trials looking to evaluate unforeseen complications of candidate haemostatic agents. Currently no generalisable assays exist for examining treatment efficacy against high titre inhibitors, and instead clinicians are forced to rely on empirical dosing and consensus guidance. With the introduction of novel agents, this landscape may be complicated further. Where data on monitoring of inhibitor treatment exist, there is little information from higher quality evidence sources to indicate how the outcomes of such practices are used to inform treatment decisions. There is a need to develop more robust evidence in this area to establish a standard of care for perioperatively monitoring haemophilia patients with inhibitors who are treated with current and emerging haemostatic therapies.

ACKNOWLEDGEMENTS

This review was funded by Roche Products Ltd and Chugai Pharma UK Ltd. The authors acknowledge Costello Medical Consulting, UK, for writing and editorial assistance (funded by Roche Products Ltd and Chugai Pharma UK Ltd).

All authors meet the International Committee of Medical Journal Editors' criteria for authorship and have made substantial contributions to the conception, design, execution or analysis and interpretation of the data:

Study conception/design and acquisition of data: DPH, CRMH, RL, GT, BDM, MM; analysis/interpretation of data: DPH, CRMH, RL, GT, BDM, MM; drafting of the publication, or revising it critically for important intellectual content: DPH, CRMH, RL, GT, BDM, MM; final approval of the publication: DPH, CRMH, RL, GT, BDM, MM.

1 CONFLICTS OF INTEREST/DISCLOSURES

2 DPH has received research support from Octapharma, Bayer and Shire, speaker
3 and/or consultancy honoraria from Pfizer, Shire, Sobi, Biomarin, Uniqure, Roche,
4 Octapharma, Novo Nordisk and Biotest; CRMH has attended advisory boards
5 organised by Roche, received research support from Novo Nordisk, Pfizer, Shire,
6 Bayer and Sobi, and acted as speaker in sponsored symposia for Pfizer, Shire, Bayer,
7 Sobi and Biotest; RL has received speaker fees from Octapharma, BPL and Bayer,
8 consultancy fees from Bayer, Octapharma, Novo Nordisk, Shire and Grifols, has
9 attended an advisory board organised by Roche, and was an investigator for the
10 HAVEN 2 study; GT is an employee of Roche Products Ltd.; BDM is an employee of
11 Costello Medical; MM has provided consultancy to CSL Behring and Novo Nordisk and
12 attended advisory boards organised by Shire and Bioverativ.

1 REFERENCES

- 2 1. Srivastava A, Brewer AK, Mauser-Bunschoten EP, et al. Guidelines for the
3 management of hemophilia. *Haemophilia*. 2013 Jan;19(1):e1-47.
- 4 2. Eckhardt CL, van Velzen AS, Peters M, et al. Factor VIII gene (F8) mutation
5 and risk of inhibitor development in nonsevere hemophilia A. *Blood*.
6 2013;122(11):1954-62.
- 7 3. Peyvandi F, Mannucci PM, Garagiola I, et al. A Randomized Trial of Factor
8 VIII and Neutralizing Antibodies in Hemophilia A. *New England Journal of Medicine*.
9 2016;374(21):2054-64.
- 10 4. Ingerslev J. Hemophilia. Strategies for the treatment of inhibitor patients.
11 *Haematologica*. 2000 Oct;85(10 Suppl):15-20.
- 12 5. Teitel JM, Carcao M, Lillicrap D, et al. Orthopaedic surgery in haemophilia
13 patients with inhibitors: a practical guide to haemostatic, surgical and rehabilitative
14 care. *Haemophilia*. 2009;15(1):227-39.
- 15 6. Collins PW, Chalmers E, Hart DP, et al. Diagnosis and treatment of factor VIII
16 and IX inhibitors in congenital haemophilia: (4th edition). *British Journal of*
17 *Haematology*. 2013;160(2):153-70.
- 18 7. Hay CRM, DiMichele DM. The principal results of the International Immune
19 Tolerance Study: a randomized dose comparison. *Blood*. 2012;119(6):1335-44.
- 20 8. van Velzen AS, Eckhardt CL, Hart DP, et al. Inhibitors in nonsevere
21 haemophilia A: outcome and eradication strategies. *Thrombosis and haemostasis*.
22 2015 //
23 22.11.2017;114(07):46-55.
- 24 9. Batty P, Austin SK, Khair K, et al. Treatment burden, haemostatic strategies
25 and real world inhibitor screening practice in non - severe haemophilia A. *British*
26 *Journal of Haematology*. 2017;176(5):796-804.
- 27 10. Ludlam CA, Smith MP, Morfini M, et al. A prospective study of recombinant
28 activated factor VII administered by continuous infusion to inhibitor patients
29 undergoing elective major orthopaedic surgery: A pharmacokinetic and efficacy
30 evaluation. *British Journal of Haematology*. 2003 01 Mar;120(5):808-13.
- 31 11. Pruthi R, Mathew P, Valentino L, Seremetis S. An open-label, randomized,
32 parallel, multi-center trial comparing the safety and efficacy of rFVIIa when
33 administered as I.V. bolus or I.V. continuous infusion to hemophilia patients with
34 Inhibitors during and after surgery. *Blood*, 2004:79b-80b.
- 35 12. Balkan C, Karapinar D, Aydogdu S, et al. Surgery in patients with haemophilia
36 and high responding inhibitors: Izmir experience. *Haemophilia*. 2010 Nov;16(6):902-
37 9.
- 38 13. Gatti L, Mannucci PM. Use of porcine factor VIII in the management of
39 seventeen patients with factor VIII antibodies. *Thrombosis and haemostasis*. 1984
40 Jul 29;51(3):379-84.
- 41 14. Kraut EH, Aledort LM, Arkin S, Stine KC, Wong WY. Surgical interventions in a
42 cohort of patients with haemophilia A and inhibitors: An experiential retrospective
43 chart review. *Haemophilia*. 2007 September;13(5):508-17.
- 44 15. Scharrer I. Recombinant factor VIIa for patients with inhibitors to factor VIII
45 or IX or factor VII deficiency. *Haemophilia*. 1999;5(4):253-9.
- 46 16. Mauser-Bunschoten EP, de Goede-Bolder A, Wielenga JJ, Levi M, Peerlinck K.
47 Continuous infusion of recombinant factor VIIa in patients with haemophilia and
48 inhibitors. Experience in The Netherlands and Belgium. *The Netherlands journal of*
49 *medicine*. 1998 Dec;53(6):249-55.

17. Mauser-Bunschoten EP, Koopman MM, Goede-Bolder AD, et al. Efficacy of recombinant factor VIIa administered by continuous infusion to haemophilia patients with inhibitors. *Haemophilia*. 2002 Sep;8(5):649-56.
18. Smith MP, Ludlam CA, Collins PW, et al. Elective surgery on factor VIII inhibitor patients using continuous infusion of recombinant activated factor VII: plasma factor VII activity of 10 IU/ml is associated with an increased incidence of bleeding. *Thrombosis and haemostasis*. 2001 Oct;86(4):949-53.
19. Pruthi RK, Mathew P, Valentino LA, et al. Haemostatic efficacy and safety of bolus and continuous infusion of recombinant factor VIIa are comparable in haemophilia patients with inhibitors undergoing major surgery. Results from an open-label, randomized, multicenter trial. *Thrombosis and haemostasis*. 2007 Oct;98(4):726-32.
20. Furukawa S, Nogami K, Ogiwara K, et al. Systematic monitoring of hemostatic management in hemophilia A patients with inhibitor in the perioperative period using rotational thromboelastometry. *Journal of Thrombosis and Haemostasis*. 2015 01 Jul;13(7):1279-84.
21. Serban M, Poenaru D, Patrascu J, et al. Risks and challenges of orthopaedic invasive interventions in haemophilia in a low-resource country. A single-center experience. *Haemostaseologie*. 2014;34 Suppl 1:S30-5.
22. Holmstrom M, Tran HT, Holme PA. Combined treatment with APCC (FEIBA(R)) and tranexamic acid in patients with haemophilia A with inhibitors and in patients with acquired haemophilia A--a two-centre experience. *Haemophilia*. 2012 Jul;18(4):544-9.
23. Habermann B, Hochmuth K, Hovy L, Scharrer I, Kurth AH. Management of haemophilic patients with inhibitors in major orthopaedic surgery by immunadsorption, substitution of factor VIII and recombinant factor VIIa (NovoSeven): a single centre experience. *Haemophilia*. 2004 Nov;10(6):705-12.
24. Danielson H, Lassila R, Ylinen P, Yrjonen T. Total joint replacement in inhibitor-positive haemophilia: Long-term outcome analysis in fifteen patients. *World j*. 2017 Oct 18;8(10):777-84.
25. Mancuso ME, Chantarangkul V, Clerici M, et al. Low thrombin generation during major orthopaedic surgery fails to predict the bleeding risk in inhibitor patients treated with bypassing agents. *Haemophilia*. 2016 Jul;22(4):e292-300.
26. Santagostino E, Mancuso ME, Novembrino C, et al. Management of Joint Replacement in Hemophilia a with Inhibitors during Emicizumab Prophylaxis. *Blood*. 2017;130(Suppl 1):2360-.
27. Kruse-Jarres R, Callaghan MU, Croteau SE, et al. Surgical experience in two multicenter, open-label phase 3 studies of emicizumab in persons with hemophilia a with inhibitors (HAVEN 1 and HAVEN 2). *Blood Conference: 59th Annual Meeting of the American Society of Hematology, ASH*. 2017;130(Supplement 1).
28. Oldenburg J, Mahlangu JN, Kim B, et al. Emicizumab Prophylaxis in Hemophilia A with Inhibitors. *The New England journal of medicine*. 2017 Aug 31;377(9):809-18.
29. Genentech. HEMLIBRA® (emicizumab-kxwh) injection, for subcutaneous use. FULL PRESCRIBING INFORMATION. https://www.gene.com/download/pdf/hemlibra_prescribing.pdf; 2017.
30. Dargaud Y, Lienhart A, Janbain M, et al. Use of thrombin generation assay to personalize treatment of breakthrough bleeds in a patient with hemophilia and inhibitors receiving prophylaxis with emicizumab. *Haematologica*. 2018.
31. Novo Nordisk A/S. A Trial Evaluating the Efficacy and Safety of Prophylactic Administration of Concizumab in Haemophilia A and B Patients With Inhibitors (explorer™4). <https://clinicaltrials.gov/ct2/show/NCT03196284>; 2017.

32. Waters EK, Sigh J, Friedrich U, Hilden I, Sorensen BB. Concizumab, an anti-tissue factor pathway inhibitor antibody, induces increased thrombin generation in plasma from haemophilia patients and healthy subjects measured by the thrombin generation assay. *Haemophilia*. 2017 Sep;23(5):769-76.
33. Pasi KJ, Rangarajan S, Georgiev P, et al. Targeting of Antithrombin in Hemophilia A or B with RNAi Therapy. *The New England journal of medicine*. 2017 Aug 31;377(9):819-28.
34. Negrier C, Ragni MV, Georgiev P, et al. Perioperative management in patients with hemophilia receiving fitusiran, an investigational RNAi therapeutic targeting antithrombin for the treatment of hemophilia. *Haemophilia*. 2018 February;24 (Supplement 1):27.
35. Collins P, Liesner R, Makris M, et al. Treatment of bleeding episodes in haemophilia A complicated by a factor VIII inhibitor in patients receiving Emicizumab. Guidance from UKHCDO Inhibitor Working Party and Executive Committee. 2018 [cited 03/05/2018]; Available from: <http://www.ukhcdo.org/wp-content/uploads/2018/01/UKHCDO-guideline-for-treatment-of-bleeds-while-on-Emicizumab-10.1.18-fi....pdf>
36. Dargaud Y, Lienhart A, Negrier C. Prospective assessment of thrombin generation test for dose monitoring of bypassing therapy in hemophilia patients with inhibitors undergoing elective surgery. *Blood*. 2010 Dec 16;116(25):5734-7.
37. Querol-Giner M, Perez-Alenda S, Iradi A, et al. Synoviorthesis in the Treatment of Recurrent Hemarthrosis in Haemophilia Patients with Inhibitors. *Haemophilia*. 2016;22:24.
38. Antmen BA, Sasmaz I, Karagun, B.S., et al. The usage of recombinant activated factor VII (RFVIIA) during major and minor surgeries in severe hemophilia patients with inhibitor. XXV Congress of the International Society on Thrombosis and Haemostasis, 2015:1–997.
39. Antmen B, Sasmaz I, Karagun B, et al. Circumcision in patients with hemophilia and the other bleeding disorders in southern part of Turkey. *Haemophilia*. 2018 February;24 (Supplement 1):61.
40. Bensadok M, Chennoukh WK, Aboura C, et al. Haemophilia with inhibitors, update from Algiers experience, about one center. XXV Congress of the International Society on Thrombosis and Haemostasis, 2015:1–997.
41. Carulli C, Rizzo AR, Linari S, et al. Joint replacements for severe haemophilic arthropathy in patients with inhibitors: A long-term experience at a single institution. *Blood Transfusion*. 2017 November;15 (Supplement 4):s546.
42. Caviglia H, Candela M, Galatro G, et al. Elective orthopaedic surgery for haemophilia patients with inhibitors: single centre experience of 40 procedures and review of the literature. *Haemophilia*. 2011 Nov;17(6):910-9.
43. Caviglia H, Galatro G, Cambiaggi G, et al. Treatment of subchondral cysts in patients with haemophilia. *Haemophilia* 2016;22:292–7.
44. Chapin J, Bamme J, Hsu F, Christos P, DeSancho M. Outcomes in Patients With Hemophilia and von Willebrand Disease Undergoing Invasive or Surgical Procedures. *Clinical and applied thrombosis/hemostasis : official journal of the International Academy of Clinical and Applied Thrombosis/Hemostasis*. 2017 Mar;23(2):148-54.
45. Ciavarella N, Antoncetti S, Ranieri P. Efficacy of porcine factor VIII in the management of haemophiliacs with inhibitors. *Br J Haematol*. 1984 Dec;58(4):641-8.
46. Dimichele D, Negrier C. A retrospective postlicensure survey of FEIBA efficacy and safety. *Haemophilia*. 2006 Jul;12(4):352-62.

- 1 47. Freiburghaus C, Berntorp E, Ekman M, et al. Immunoabsorption for removal
2 of inhibitors: update on treatments in Malmö-Lund between 1980 and 1995.
3 *Haemophilia*. 1998 Jan;4(1):16-20.
- 4 48. He Y, Zhou X, Cui H, et al. Surgical Management of Haemophilic
5 Pseudotumors: Experience in a Developing Country. *Journal of Investigative Surgery*.
6 2017 31 Oct:1-10.
- 7 49. Ingerslev J, Freidman D, Gastineau D, et al. Major surgery in haemophilic
8 patients with inhibitors using recombinant factor VIIa. *Haemostasis*. 1996;26 Suppl
9 1:118-23.
- 10 50. Ingerslev J. Efficacy and safety of recombinant factor VIIa in the prophylaxis
11 of bleeding in various surgical procedures in hemophilic patients with factor VIII and
12 factor IX inhibitors. *Seminars in thrombosis and hemostasis*. 2000;26(4):425-32.
- 13 51. Jenkins PJ, Ekrol I, Lawson GM. Total knee replacement in patients with
14 haemophilia: the Scottish experience. *Scottish medical journal*. 2013 Nov;58(4):223-
15 7.
- 16 52. Kitchens CS. Surgery in hemophilia and related disorders. A prospective study
17 of 100 consecutive procedures. *Medicine*. 1986 Jan;65(1):34-45.
- 18 53. Kizilocak H, Ozdemir N, Ozcan R, Celkan T. Circumcision in children with
19 haemophilia. *Journal of Thrombosis and Haemostasis*. 2016;14(Suppl 1):1-168.
- 20 54. Lauroua P, Ferrer AM, GuÉRin V. Successful major and minor surgery using
21 factor VIII inhibitor bypassing activity in patients with haemophilia A and inhibitors.
22 *Haemophilia*. 2009;15(6):1300-7.
- 23 55. Lim MY, Nielsen B, Lee K, et al. Rituximab as first-line treatment for the
24 management of adult patients with non-severe hemophilia A and inhibitors. *Journal*
25 *of thrombosis and haemostasis : JTH*. 2014 Jun;12(6):897-901.
- 26 56. Lozier JN, Santagostino E, Kasper CK, Teitel JM, Hay CR. Use of porcine
27 factor VIII for surgical procedures in hemophilia A patients with inhibitors. *Seminars*
28 *in hematology*. 1993 Apr;30(2 Suppl 1):10-21.
- 29 57. Mahasandana C, Patharathienskul D, Suvatte V. Hemophilia with factor VIII
30 and factor IX inhibitors, incidence, bleeding problems and management. *The*
31 *Southeast Asian journal of tropical medicine and public health*. 1993;24 Suppl 1:106-
32 12.
- 33 58. Mancuso ME, Mannucci PM, Sartori A, Agliardi A, Santagostino E. Feasibility of
34 prophylaxis and immune tolerance induction regimens in haemophilic children using
35 fully implantable central venous catheters. *Br J Haematol*. 2008 May;141(5):689-95.
- 36 59. Morado M, Jimenez-Yuste V, Villar A, et al. Complications of central venous
37 catheters in patients with haemophilia and inhibitors. *Haemophilia*. 2001
38 Nov;7(6):551-6.
- 39 60. Nguyen T, Tran D, Dao H, et al. Surgical interventions in patients with
40 hemophilia: Outcome of treatment at a single center in low-resource settings.
41 *Haemophilia*. 2018 February;24 (Supplement 1):130.
- 42 61. Nilsson IM, Hedner U, Ahlberg A, Larsson SA, Bergentz SE. Surgery of
43 hemophiliacs--20 years' experience. *World journal of surgery*. 1977 Jan;1(1):55-66.
- 44 62. O'Connell N, Mc Mahon C, Smith J, et al. Recombinant factor VIIa in the
45 management of surgery and acute bleeding episodes in children with haemophilia
46 and high responding inhibitors. *Br J Haematol*. 2002 Mar;116(3):632-5.
- 47 63. Oldenburg J, Shima M, Kruse-Jarres R, et al. Bleeding events and safety
48 outcomes in pediatric persons with hemophilia a with inhibitors: The first non-
49 interventional study (NIS) from a real-world setting. *Blood Conference: 59th Annual*
50 *Meeting of the American Society of Hematology, ASH*. 2017;130(Supplement 1).
- 51 64. Özdemir GN, Çelik E, Bulut M, et al. Circumcision in children with bleeding
52 diathesis. *Turkish Archives of Pediatrics*. 2011;46:304-8.

65. Quintana-Molina M, Martinez-Bahamonde F, Gonzalez-Garcia E, et al. Surgery in haemophilic patients with inhibitor: 20 years of experience. *Haemophilia*. 2004 Sep;10 Suppl 2:30-40.
66. Rodriguez-Merchan EC. Total knee replacement in haemophilic arthropathy. *The Journal of bone and joint surgery British volume*. 2007 Feb;89(2):186-8.
67. Rodriguez-Merchan EC, Jimenez-Yuste V, Gomez-Cardero P, et al. Surgery in haemophilia patients with inhibitors, with special emphasis on orthopaedics: Madrid experience. *Haemophilia*. 2010 May;16(102):84-8.
68. Sasmaz I, Antmen B, Leblebisatan G, et al. Circumcision and complications in patients with haemophilia in southern part of Turkey: Cukurova experience. *Haemophilia*. 2012 May;18(3):426-30.
69. Sasmaz I, Antmen BA, Karagun BS, et al. The experience of surgery with activated prothombin complex concentrates in haemophilia a patients with inhibitors. *Haemophilia*. 2015;21(Suppl 2):14–94.
70. Sasmaz I, Antmen B, Karagun B, et al. The surgeries in hemophilia A patients with inhibitors. *Haemophilia*. 2018 February;24 (Supplement 1):84-5.
71. Szczepanik AB, Wislawski S, Windyga J, et al. [Strategy for secure hemostasis in hemophilia patients undergoing surgery for malignant neoplasms]. *Polski merkuriusz lekarski : organ Polskiego Towarzystwa Lekarskiego*. 2009 Nov;27(161):375-80.
72. Serban M, Poenaru D, Pop L, et al. Surgery--a challenge in haemophiliacs with inhibitors. *Hamostaseologie*. 2009 Oct;29 Suppl 1:S39-41.
73. Shapiro AD, Gilchrist GS, Hoots WK, Cooper HA, Gastineau DA. Prospective, randomised trial of two doses of rFVIIa (NovoSeven) in haemophilia patients with inhibitors undergoing surgery. *Thrombosis and haemostasis*. 1998 Nov;80(5):773-8.
74. Shapiro A, Cooper DL. U.S. survey of surgical capabilities and experience with surgical procedures in patients with congenital haemophilia with inhibitors. *Haemophilia*. 2012 May;18(3):400-5.
75. Smith OP. Recombinant factor VIIa in the management of surgery and acute bleeding episodes in children with haemophilia and high-responding inhibitors. *Pathophysiology of haemostasis and thrombosis*. 2002;32 Suppl 1:22-5.
76. Solimeno LP, Mancuso ME, Pasta G, et al. Factors influencing the long-term outcome of primary total knee replacement in haemophiliacs: a review of 116 procedures at a single institution. *Br J Haematol*. 2009 Apr;145(2):227-34.
77. Takedani H, Shima M, Horikoshi Y, et al. Ten-year experience of recombinant activated factor VII use in surgical patients with congenital haemophilia with inhibitors or acquired haemophilia in Japan. *Haemophilia*. 2015 May;21(3):374-9.
78. Gringeri A, Fischer K, Karafoulidou A, et al. Sequential combined bypassing therapy is safe and effective in the treatment of unresponsive bleeding in adults and children with haemophilia and inhibitors. *Haemophilia*. 2011 Jul;17(4):630-5.
79. Lak M, Sharifian RA, Karimi K, Mansouritorghabeh H. Acquired hemophilia A: clinical features, surgery and treatment of 34 cases, and experience of using recombinant factor VIIa. *Clinical and applied thrombosis/hemostasis : official journal of the International Academy of Clinical and Applied Thrombosis/Hemostasis*. 2010 Jun;16(3):294-300.
80. Liozon E, Delaire L, Turlure P. Acquired inhibitor to factor VIII: C in non hemophiliacs (acquired hemophilia). Clinico-biologic study and management in nine patients. [French]. *Annales de Medecine Interne* 1997;148:477–90.
81. Ma AD, Kessler CM, Al-Mondhiry HA, Gut RZ, Cooper DL. US experience with recombinant factor VIIa for surgery and other invasive procedures in acquired haemophilia: analysis from the Hemostasis and Thrombosis Research Society Registry. *Haemophilia*. 2016 Jan;22(1):e18-24.

82. Novack A, St-Louis J, Greist A, et al. Perioperative management of bleeds with recombinant porcine FVIII in patients with acquired hemophilia A. *Journal of Thrombosis and Haemostasis*. 2015;13(Suppl 2):1997.
83. Zulfikar B, Kilicoglu O, Atalar A, et al. Hemophilic Arthropathy in Inhibitor Positive Hemophilia Patients. *Haemophilia*. 2016;22:85-6.
84. Boadas A, Fernandez-Palazzi F, De Bosch NB, Cedeno M, Ruiz-Saez A. Elective surgery in patients with congenital coagulopathies and inhibitors: experience of the National Haemophilia Centre of Venezuela. *Haemophilia*. 2011 May;17(3):422-7.
85. Carulli C, Rizzo AR, Martini C, et al. The orthopaedic treatment of haemophilic arthropathy in patients with inhibitors: A 15-year experience at a single Institution. *Haemophilia*. 2016;22(Suppl. 4):23.
86. Castaman G, Linari S, Carulli C, Innocenti M. Short and long-term outcomes in hemophilia patients with inhibitors undergoing orthopedic prosthetic surgery. *Blood*. 2015;126(23):4704.
87. Croteau SE, Nakar C, Neufeld EJ, Shapiro A, Cooper DL. Safety and efficacy of recombinant factor VIIa by pediatric age cohort: reassessment of compassionate use and trial data supporting US label. *Pediatric blood & cancer*. 2016 Oct;63(10):1822-8.
88. Hilgartner MW, Knatterud GL. The use of factor eight inhibitor by-passing activity (FEIBA immuno) product for treatment of bleeding episodes in hemophiliacs with inhibitors. *Blood*. 1983 Jan;61(1):36-40.
89. Ju HY, Jang HL, Park YS. The efficacy of bypassing agents in surgery of hemophilia patients with inhibitors. *Blood research*. 2015 Sep;50(3):173-8.
90. Kavakli K, Zulfikar H, Zulfikar B, et al. Efficacy of FEIBA for acute bleeding and surgical haemostasis in haemophilia A patients with inhibitors: A multicentre registry in Turkey. *Haemophilia*. 2012 May;18(3):383-91.
91. Linari S, Carulli C, Martini C. Short and long-term outcomes in hemophilia patients with inhibitors undergoing orthopedic prosthetic surgery. *Haematologica*. 2015;100:26-7.
92. Negrier C, Goudemand J, Sultan Y, et al. Multicenter retrospective study on the utilization of FEIBA in France in patients with factor VIII and factor IX inhibitors. French FEIBA Study Group. Factor Eight Bypassing Activity. *Thrombosis and haemostasis*. 1997 Jun;77(6):1113-9.
93. Negrier C, Lienhart A, Numerof R, et al. SURgical interventions with FEIBA (SURF): international registry of surgery in haemophilia patients with inhibitory antibodies. *Haemophilia*. 2013 May;19(3):e143-50.
94. Polyanskaya T, Zorenko V, Karpov E, et al. Experience of recombinant activated factor VII usage during surgery in patients with haemophilia with inhibitors. *Haemophilia*. 2012 Nov;18(6):997-1002.
95. Rangarajan S, Yee TT, Wilde J. Experience of four UK comprehensive care centres using FEIBA(R) for surgeries in patients with inhibitors. *Haemophilia*. 2011 Jan;17(1):28-34.
96. Rodriguez-Merchan EC. Surgery in haemophilic patients with inhibitors. *Haemophilia*. 2004 Sep;10 Suppl 2:1-2.
97. Santagostino E, Morfini M, Rocino A, et al. Relationship between factor VII activity and clinical efficacy of recombinant factor VIIa given by continuous infusion to patients with factor VIII inhibitors. *Thrombosis and haemostasis*. 2001 Oct;86(4):954-8.
98. Sasmaz I, Antmen B, Guvenc B, et al. Circumcision and complications in adolescent and adult patients with hemophilia in southern part of Turkey. 20th Congress Of The European Hematology Association. Vienna, Austria, 2015:1-804.

- 1 99. Scharf R, Kucharski W, Nowak T. Surgery in hemophilia A patients with factor
2 VIII inhibitor: 10-year experience. *World journal of surgery*. 1996 Nov-
3 Dec;20(9):1171-81.
- 4 100. Tjonnfjord GE. Activated prothrombin complex concentrate (FEIBA) treatment
5 during surgery in patients with inhibitors to FVIII/IX: the updated Norwegian
6 experience. *Haemophilia*. 2004 Sep;10 Suppl 2:41-5.
- 7 101. Tjonnfjord GE. Surgery in patients with hemophilia and inhibitors: a review of
8 the Norwegian experience with FEIBA. *Seminars in hematology*. 2006 Apr;43(2 Suppl
9 4):S18-21.

10

1 **FIGURE LEDGEND**

2 **Figure 1. Systematic Review of Clinical Studies**

3 [†]Only articles discussing laboratory monitoring were extracted




Table 1. Studies Included Following Full Text Screening

Study	Study Type	Country	Monitoring reported
Congenital haemophilia			
Alenda et al. 2016[37]	Observational	Spain	Not reported
Antmen et al. 2015[38]	Observational	Turkey	Not reported
Antmen et al. 2018[39]	Observational	Turkey	Not reported
Balkan et al. 2010[12]	Observational	Turkey	Laboratory monitoring
Bensadok et al. 2015[40]	Observational	Algeria	Not reported
Carulli et al. 2017[41]	Observational	Italy	Not reported
Caviglia et al. 2011[42]	Observational	Argentina	Not reported
Caviglia et al. 2016[43]	Observational	Argentina	Not reported
Chapin et al. 2017[44]	Observational	USA	Not reported
Ciavarella et al. 1984[45]	Interventional	Italy	Not reported
Danielson et al. 2017[24]	Observational	Finland	Laboratory monitoring
Dimichele et al. 2006[46]	Observational	USA and Europe	Not reported
Freiburghaus et al. 1998[47]	Observational	Sweden	Not reported
He et al. 2017[48]	Observational	China	Not reported
Holmström et al. 2012[22]	Interventional	Norway and Sweden	Laboratory monitoring
Ingerslev et al. 1996[49]	Observational	Multiple	Laboratory monitoring
Ingerslev et al. 2000[50]	Observational	Denmark	Not reported
Jenkins et al. 2013[51]	Observational	UK	Not reported
Karagun et al. 2016	Observational	Turkey	Not reported
Kitchens et al. 1986[52]	Observational	USA	Not reported
Kizilocak et al. 2016[53]	Observational	Turkey	Not reported
Kruse-Jarres et al. 2017[27]	Interventional	Multiple	Not reported
Lauroua et al. 2009[54]	Observational	France	Both [clinical and laboratory monitoring]
Lim et al. 2014[55]	Observational	USA	Not reported

Study	Study Type	Country	Monitoring reported
Lozier et al. 1993[56]	Observational	Multiple	Not reported
Ludlam et al. 2003[10]	Interventional	UK and Italy	Laboratory monitoring
Mahasandana et al. 1993[57]	Observational	Thailand	Not reported
Mancuso et al. 2008[58]	Observational	Italy	Not reported
Morado et al. 2001[59]	Observational	Spain	Not reported
Negrier et al. 2018[34]	Interventional	Multiple	Not reported
Nguyen et al. 2018[60]	Observational	Vietnam	Not reported
Nilsson et al. 1977[61]	Observational	Sweden	Laboratory monitoring†
O'Connell et al. 2002[62]	Observational	Ireland and UK	Clinical monitoring
Oldenburg et al. 2017[63]	Observational	Multiple	Not reported
Ozdemir et al. 2011[64]	Observational	Turkey	Not reported
Pruthi et al. 2007[19]	Interventional	USA	Laboratory monitoring
Quintana-Molina et al. 2004[65]	Observational	Spain	Laboratory monitoring
Rodriguez-Merchan et al. 2007[66]	Observational	Spain	Not reported
Rodriguez-Merchan et al. 2010[67]	Observational	Spain	Not reported
Sasmaz et al. 2012[68]	Observational	Turkey	Not reported
Sasmaz et al. 2015[69]	Observational	Turkey	Not reported
Sasmaz et al. 2018[70]	Observational	Turkey	Not reported
Szczepanik et al. 2018[71]	Observational	Poland	Not reported
Serban et al. 2009[72]	Observational	Romania	Not reported
Shapiro et al. 1998[73]	Interventional	USA	Laboratory monitoring
Shapiro et al. 2012[74]	Observational	USA	Not reported
Smith et al. 2002[75]	Observational	Ireland and UK	Both [clinical and laboratory monitoring]

Study	Study Type	Country	Monitoring reported
Solimeno et al. 2009[76]	Observational	Italy	Not reported
Takedani et al. 2010[77]	Observational	Japan	Not reported
Acquired haemophilia			
Gringeri et al. 2011[78]	Observational	Europe	Laboratory monitoring
Lak et al. 2010[79]	Observational	Iran	Laboratory monitoring
Liozon et al. 1997[80]	Observational	France	Laboratory monitoring
Ma et al. 2016[81]	Interventional	USA	Not reported
Novack et al. 2015[82]	Observational	Multiple	Not reported
Both congenital and acquired haemophilia			
Atalar et al. 2016[83]	Observational	Turkey	Not reported
Boadas et al. 2011[84]	Observational	Venezuela	Both [clinical and laboratory monitoring] ⁺
Carulli et al. 2016[85]	Observational	Italy	Not reported
Castaman et al. 2015[86]	Observational	Italy	Not reported
Croteau et al. 2016[87]	Interventional	USA	Not reported
Furukawa et al. 2015[20]	Interventional	Japan	Laboratory monitoring
Gatti et al. 1984[13]	Interventional	Italy	Laboratory monitoring
Habermann et al. 2004[23]	Observational	Germany	Laboratory monitoring
Hilgartner et al. 1983[88]	Observational	USA	Not reported
Ju et al. 2015[89]	Observational	South Korea	Clinical monitoring
Kavakli et al. 2012[90]	Observational	Turkey	Not reported
Kraut et al. 2007[14]	Observational	USA	Laboratory monitoring
Linari et al. 2015[91]	Observational	Italy	Not reported
Mancuso et al. 2016[25]	Observational	Italy	Laboratory monitoring
Mauser-Bunschoten et al. 1998[16]	Observational	Netherlands and Belgium	Both [clinical and laboratory monitoring]
Mauser-Bunschoten et al. 2002[17]	Observational	Netherlands	Both [clinical and laboratory monitoring]
Negrier et al. 1997[92]	Observational	France	Not reported

Study	Study Type	Country	Monitoring reported
Negrier et al. 2013[93]	Observational	Worldwide: Colombia, France, Germany, Italy, South Korea, Sweden and the UK	Both [clinical and laboratory monitoring]
Polyanskaya et al. 2012[94]	Observational	Russia	Clinical monitoring
Rangarajan et al. 2011[95]	Observational	UK	Clinical monitoring
Rodriguez-Merchan et al. 2004[96]	Observational	Worldwide	Not reported
Rodriguez-Merchan et al. 2007[66]	Observational	Spain	Not reported
Santagostino et al. 2001[97]	Observational	Italy	Laboratory monitoring
Sasmaz et al. 2015[98]	Observational	Turkey	Not reported
Scharf et al. 1996[99]	Observational	Poland	Not reported
Scharrer et al. 1999[15]	Interventional	Germany	Both [clinical and laboratory monitoring]
Serban et al. 2014[21]	Observational	Romania	Both [clinical and laboratory monitoring]
Smith et al. 2001[18]	Interventional	Unclear	Both [clinical and laboratory monitoring]
Szczepanik et al. 2009[71]	Observational	Poland	Not reported
Takedani et al. 2015[77]	Observational	Japan	Clinical monitoring
Tjonnfjord et al. 2004/ Tjonnfjord et al. 2006[100, 101]	Observational	Norway	Laboratory monitoring

[†]Outcomes not reported separately for inhibitor patients so data not extracted; [‡]No monitoring results reported so data not extracted

Table 2. Overview of Studies Reporting Perioperative Laboratory Monitoring

Study	Study design	Patients	Procedures	Haemostatic treatment	Haemostatic outcome
Balkan C et al. 2010[12]	Single-centre, retrospective observational study	30 HA patients with high responding inhibitors	11 major 42 minor	<ul style="list-style-type: none"> • aPCC, or • rFVIIa, or • Sequential use of aPCC and rFVIIa 	<ul style="list-style-type: none"> • aPCC: 22/22 (100%) bleeding controlled • rFVIIa: 31/33 (94%) bleeding controlled
Danielson H et al. 2017[24]	Single-centre, retrospective observational study	6 HA patients with inhibitors (n=2 low-responding, n=4 high-responding)	15 orthopaedic	<ul style="list-style-type: none"> • Cryoprecipitate, or • Coagulation FVIII (pdFVIII or rFVIII), or • aPCC, or • rFVIIa (post-treatment switch in some individual cases) 	8/15 (53%) bleeding controlled (rated as 'good', indicating no difference in bleeding compared to normal arthroplasty)
Furukawa S et al. 2015[20]	Single-centre, prospective interventional study	8 HA patients with inhibitors	8 elective	<ul style="list-style-type: none"> • rFVIIa, or • aPCC 	8/8 (100%) bleeding controlled
Gatti L et al. 1984[13]	Single-centre, prospective, uncontrolled interventional study	5 HA patients with inhibitors	3 minor 2 major	<ul style="list-style-type: none"> • Bolus porcine FVIII (Hyate:C) (minor dental), or • Continuous porcine FVIII (Hyate:C) (major) 	2/2 (100%) bleeding controlled (only reported for major surgery)
Habermann B et al. 2004[23]	Single-centre, retrospective observational study	4 HA patients with inhibitors	6 orthopaedic	Anvitoff™ (containing TXA) in combination with: <ul style="list-style-type: none"> • bolus FVIII (low inhibitor titre) • immunoabsorbant therapy (Therasorb™) followed by bolus FVIII (high inhibitor titre) • continuous rFVIIa infusion when inhibitor titres rose/could not be eliminated or FVIII response decreased 	5/6 (83%) bleeding controlled
Holmström M et al. 2012[22]	Two-centre, prospective interventional study	6 HA patients with high responding inhibitors	2 minor 5 major	Bolus aPCC in combination with TXA	6/7 (86%) bleeding controlled

Study	Study design	Patients	Procedures	Haemostatic treatment	Haemostatic outcome
Ingerslev J. et al. 1996[49]	Multicentre, retrospective observational study	11 HA patients and 1 HB patient with inhibitors	13 major	Bolus rFVIIa	12/12 (100%) bleeding controlled (Outcome not reported in n=1 case)
Kraut EH et al. 2007[14]	Multicentre, retrospective chart review	6 HA patients with inhibitors	21 various	<ul style="list-style-type: none"> • Bolus aPCC monotherapy, or • Bolus rFVIIa monotherapy, or • Bolus/continuous combination therapy 	14/21 (67%) bleeding controlled
Lauroua P et al. 2009[54]	Single-centre, retrospective observational study	7 HA patients with inhibitors	8 major elective 2 major emergency 2 minor elective	Bolus aPCC as first-line treatment	Haemostatic outcomes were consistent with non-coagulopathic patients undergoing similar procedures
Ludlam A et al. 2003[10]	Prospective, interventional study	9 HA patients with inhibitors	9 major orthopaedic	Continuous rFVIIa	8/9 (88.9%) bleeding controlled at end of surgery
Mancuso ME et al. 2016[25]	Single-centre, prospective, observational study	10 HA patients with inhibitors	11 major orthopaedic	Bolus doses of: <ul style="list-style-type: none"> • rFVIIa • aPCC • Sequential therapy with rFVIIa and aPCC 	10/11 (91%) bleeding controlled
Mauser-Bunschoten EP et al. 1998[16]	Multicentre, retrospective observational study	3 HA patients with inhibitors	2 dental extraction 2 hip arthroplasty	Continuous rFVIIa	3/4 (75%) bleeding controlled
Mauser-Bunschoten EP et al. 2002[17]	Multicentre, prospective observational study	4 HA patients and 1 HB patient with inhibitors	2 synovectomy 4 dental extraction 1 orthopaedic surgery	Continuous rFVIIa	NR (except 2 dental extractions rated 'ineffective' and 2 rated 'partially effective')
Négrier C et al. 2013[93]	Multicentre, prospective, observational study	18 HA patients and 2 HB patients with inhibitors	35 various (including procedures performed on n=4 acquired haemophilia patients)	Bolus aPCC	31/34 (91%) bleeding controlled (rated as 'excellent' or 'good'; full population including acquired haemophilia patients; concomitant medication documented in n=34/35 surgical procedures)
Pruthi RK et al. 2007[19]	Multicentre, prospective interventional study	24 HA/HB patients with inhibitors (A/B subgroups not specified)	24 elective surgery	Bolus and continuous infusion of rFVIIa	17/23 (74%) bleeding controlled overall (n=1 patient excluded from efficacy analysis)

Study	Study design	Patients	Procedures	Haemostatic treatment	Haemostatic outcome
Quintana-Molina M et al. 2004[65]	Single-centre, retrospective observational study	45 HA patients 3 HB patients with inhibitors	10 major elective and emergency 54 minor elective and emergency	Bolus doses of: <ul style="list-style-type: none"> • rFVIIa, or • aPCC, or • FVIII concentrate 	<ul style="list-style-type: none"> • rFVIIa: 14/18 (78%) bleeding controlled • aPCC: 31/32 (97%) bleeding controlled • FVIII concentrate: 15/15 (100%) bleeding controlled (based on outcomes reported in article tables)
Santagostino E et al. 2001[97]	Multicentre, prospective, observational study	25 HA patients with inhibitors (unclear how many had surgery)	11 major 14 minor	Continuous rFVIIa	Surgical patients' results not reported separately
Scharrer I. 1999[15]	Multicentre, prospective interventional study	19 HA/HB patients with inhibitors (A/B subgroups not specified)	5 major 17 minor (full population including patients with acquired inhibitors/FVII deficiency)	Bolus rFVIIa	100% minor/60% major surgical procedures bleeding controlled (during surgery)
Serban M et al. 2014[21]	Single-centre, retrospective observational study	13 HA/B patients with inhibitors (not clear whether A or B)	Invasive orthopaedic (n NR)	Bolus doses and continuous infusion of: <ul style="list-style-type: none"> • FVIII/FIX concentrates • rFVIIa 	Reported but not for population of interest
Shapiro AD et al. 1998[73]	Multicentre, prospective interventional study	25 HA patients and 3 HB patients with inhibitors	29 (including 1 procedure for a patient with acquired haemophilia): 11 major 18 minor	Bolus rFVIIa	23/29 (79%) bleeding controlled (may include 1 acquired haemophilia patient's procedure)
Smith MP et al. 2001[18]	Multicentre, prospective interventional study	6 HA patients with inhibitors	6 major	Bolus dose followed by continuous infusion of rFVIIa	2/6 bleeding controlled
Smith OP et al. 2002[75]	Two-centre, retrospective chart review	12 HA patients with inhibitors	19 CVAD insertion/removal 1 multiple dental extraction	Bolus rFVIIa	20/20 (100%) bleeding controlled (n=2 cases of minor bleeding after treatment had ended were resolved with re-treatment of rFVIIa)
Tjonnfjord GE. 2004, 2006[100, 101]	Single-centre, retrospective observational study	8 HA patients with inhibitors	12 minor 6 major	Bolus aPCC	18/18 (100%) bleeding controlled

aPCC: Activated prothrombin complex concentrate; CVAD: Central venous access device; FIX: Factor IX; FVII: Factor VII; FVIII: Factor VIII; HA: Haemophilia A; HB: Haemophilia B; NR: Not reported; pdFVIII: Plasma-derived factor VIII; rFVIIa: Recombinant factor VIIa; rFVIII: Recombinant factor VIII; TXA: Tranexamic acid

Table 3: Routine Laboratory Testing

Study	Monitoring methods	Monitoring results
Balkan C et al. 2010[12]	Laboratory assessment (post-operative) of: <ul style="list-style-type: none"> • Platelet count • PT • APTT • Fibrinogen • D-dimer 	<ul style="list-style-type: none"> • APTT did not return to normal by using the haemostatic agents • Significant shortening of PT
Gatti L et al. 1984[13]	Laboratory assessment of: <ul style="list-style-type: none"> • Clinical effectiveness • The prevalence of anamnestic antibody responses and of severe or milder side effects • Platelet counts • Haematocrit 	NR
Habermann B et al. 2004[23]	Laboratory assessment of: <ul style="list-style-type: none"> • D-dimer 	<ul style="list-style-type: none"> • Only on the day of surgery was a slight increase of the D-dimer level seen. On the postoperative days, the D-dimer levels were within the normal range.
Ingerslev J. et al. 1996[49]	Laboratory assessment of: <ul style="list-style-type: none"> • Platelet count • PT • APTT • Fibrinogen • D-dimer • ATIII 	<ul style="list-style-type: none"> • Small reductions in platelet numbers • Significantly shortened PT following infusion • APTT shortened in nearly all patients • Insignificant changes in fibrinogen • All but one D-dimer sample showed results below the limits of specified abnormality • ATIII showed no tendency to decrease
Kraut EH et al. 2007[14]	Laboratory assessment, including platelet function analysis, of: <ul style="list-style-type: none"> • D-dimer levels • Haemoglobin level • Aggregation 	<ul style="list-style-type: none"> • aPCC treatment was reduced after monitoring indicated an elevation in D-dimer levels
Lauroua P et al. 2009[54]	Consumption coagulopathy and thrombogenicity evaluated with laboratory assessment of: <ul style="list-style-type: none"> • Platelets • Fibrinogen • D-dimer or fibrinogen and fibrin degradation products • Haemoglobin level 	<ul style="list-style-type: none"> • Monitoring of D-dimer, fibrinogen and fibrin degradation products showed no consistent activation of coagulation or increase in fibrinolysis • Neither platelet consumption nor fibrinogen depletion observed post-operatively • Haemoglobin remained stable above 8 g/dL in most cases
Ludlam A et al. 2003[10]	Laboratory assessment of: <ul style="list-style-type: none"> • Complete blood counts • Fibrinogen • D-dimer • ATIII (assessed by chromogenic determination) 	<ul style="list-style-type: none"> • ATIII, fibrinogen and platelet counts fluctuated but did not decline progressively • During the first 72 h of infusion, mean platelet count decreased; Mean ATIII decreased between wound closure and at 72 h

Study	Monitoring methods	Monitoring results
Mancuso ME et al. 2016[25]	Laboratory assessment of: <ul style="list-style-type: none"> Fibrinogen (Functional Clauss method) D-dimer (Latex enhanced turbidimetric immunoassay) PT (PT-based one-stage assay) 	<ul style="list-style-type: none"> D-dimer significantly increased over the first four post-operative days Fibrinogen slightly decreased on the first post-operative day, then increased for the following three post-operative days PT increased slightly in aPCC-treated patients over four post-operative days
Mauser-Bunschoten EP et al. 1998[16]	Laboratory assessment of: <ul style="list-style-type: none"> PT 	NR
Mauser-Bunschoten EP et al. 2002[17]	Laboratory assessment of: <ul style="list-style-type: none"> PT 	NR
Négrier C et al. 2013[93]	Laboratory assessment of: <ul style="list-style-type: none"> Haemoglobin Red blood cell count Haematocrit Liver enzyme levels 	<ul style="list-style-type: none"> Abnormal, significant haemoglobin levels were observed in 5 patients with inhibitors
Pruthi RK et al. 2007[19]	Laboratory assessment of: <ul style="list-style-type: none"> Fibrinogen D-dimer PT 	<ul style="list-style-type: none"> No statistically significant differences between pre- and postoperative platelet counts, fibrinogen, D-dimer and F 1.2 concentrations between bolus infusion, continuous infusion or control subjects
Quintana-Molina M et al. 2004[65]	Laboratory assessment (postoperative and control tests at least every 48 hours) of: <ul style="list-style-type: none"> Platelet count (obtained by impedance and optically) PT and cephaline time (monitored by two apparatuses based on different techniques, either optical density or magnetic force) Fibrinogen (Clauss method) D-dimer (turbidimetry) 	NR
Santagostino E et al. 2001[97]	Laboratory assessment of: <ul style="list-style-type: none"> PT APTT Fibrinogen D-dimer Platelet count 	<ul style="list-style-type: none"> Platelet count decreased during 2 courses of treatment given for knee replacement
Scharrer I. 1999[15]	Laboratory assessment of PT and APTT plus: <ul style="list-style-type: none"> Fibrinogen, or Platelets, or 	NR (Laboratory results only collected if considered necessary by the investigator and adverse event had occurred)

Study	Monitoring methods	Monitoring results
	<ul style="list-style-type: none"> • Thrombin-antithrombin complex, or • D-dimer, or • Fibrino-peptide A, or • Fibrin-degradation products, or • Fibrin monomer, or • ATIII, or • α-antiplasmin 	
Shapiro AD et al. 1998[73]	Laboratory assessment of: <ul style="list-style-type: none"> • PT • Fibrinogen • D-dimer • ATIII • Platelet count 	<ul style="list-style-type: none"> • Mean PT decreased • D-dimer levels increased in 83% of patients during the first 48 h postoperatively • No changes in ATIII • Mean fibrinogen levels increased • No change in platelet levels
Smith MP et al. 2001[18]	Laboratory assessment of: <ul style="list-style-type: none"> • International normalised ratio • Fibrinogen (Clauss method) • ATIII • Automated full blood counts • D-dimer 	<ul style="list-style-type: none"> • ATIII, fibrinogen and platelet counts were not observed to decline
Smith OP et al. 2002[75]	Laboratory assessment of: <ul style="list-style-type: none"> • PT levels 	<ul style="list-style-type: none"> • PT shortened to lower limit of normal following rFVIIa treatment • Some haemoglobin levels dropped below 8 g/dL in patients who experienced bleeding episodes
Tjonnfjord GE. 2004, 2006[100, 101]	Laboratory assessment of: <ul style="list-style-type: none"> • PT • APTT • Fibrinogen • D-dimer 	PT shortened

aPCC: Activated prothrombin complex concentrate; APTT: Activated partial thromboplastin time; ATIII: Antithrombin; NR: Not reported; PT: Prothrombin time

Table 4: Factor VII Monitoring

Study	Haemostatic treatment	Reported factors monitored	Method of monitoring	Monitoring results
Ingerslev J. et al. 1996[49]	Bolus rFVIIa	Levels of post-infusion FVII:C	Laboratory assessment	NR
Ludlam A et al. 2003[10]	Continuous rFVIIa	<ul style="list-style-type: none"> FVII:C FVIIa:C levels 	<ul style="list-style-type: none"> Plasma FVII:C was assessed by an automated one-stage FVII clot method on an automated laboratory analyser FVIIa:C was assessed using a specific automated assay 	<ul style="list-style-type: none"> Mean (range) FVII:C levels: <ul style="list-style-type: none"> Effective haemostasis, end of surgery: 37 IU/ml, (29–51 IU/ml), n=8 Ineffective haemostasis, end of surgery: 27 IU/ml, n=1 Effective haemostasis, 8h after wound closure: 38 IU/ml (24–79 IU/ml), n=5 Partially effective haemostasis, 8h after wound closure: 42 IU/ml, (37–57 IU/ml), n=4 Mean (range) FVIIa:C levels: <ul style="list-style-type: none"> Effective haemostasis, end of surgery: 50 IU/ml, (37–59 IU/ml), n=8 Ineffective haemostasis, end of surgery: 40 IU/ml, n=1 Effective haemostasis, 8h after wound closure: 52 IU/ml (37–74 IU/ml), n=5 Partially effective haemostasis, 8h after wound closure: 61 IU/ml, (38–82 IU/ml), n=4 FVII:C was >30 IU/ml at time of all but one bleeds
Mausser-Bunschoten EP et al. 1998[16]	Continuous rFVIIa	Plasma FVIIa levels	FVIIa: one-stage coagulation assay	<ul style="list-style-type: none"> FVIIa levels maintained above 10 U/ml through flow rate adjustment
Mausser-Bunschoten EP et al. 2002[17]	Continuous rFVIIa	Plasma FVIIa levels	FVIIa: one-stage coagulation assay	NR
Pruthi RK et al. 2007[19]	Bolus and continuous infusion of rFVIIa	FVII:C	Samples for FVII:C were collected within 30 min prior to and at 10 min after the initial 90 µg/kg rFVIIa bolus infusion, at 0, 8, 24, 48 and 72 h after wound closure and daily from post-operative day 4 –10 or until discharge (and prior to any supplemental bolus	<ul style="list-style-type: none"> At wound closure, FVII:C levels were higher in continuous vs. bolus infusion patients, which as sustained through 72 h but not statistically significant In subjects for whom therapy was ineffective, FVII:C levels were in excess of 30 IU/ml at the time therapy was declared ineffective

Study	Haemostatic treatment	Reported factors monitored	Method of monitoring	Monitoring results
			infusion of rFVIIa). FVII:C was measured in a central laboratory	
Santagostino E et al. 2001[97]	Continuous rFVIIa	FVII:C	One-stage coagulation assay	<ul style="list-style-type: none"> FVII:C levels were significantly higher during continuous infusion courses given for major surgery than minor surgery rFVIIa clearance was significantly lower in courses given for major surgery than for minor surgery
Shapiro AD et al. 1998[73]	Bolus rFVIIa	FVII:C	Laboratory analysis of blood sample	<ul style="list-style-type: none"> FVII:C could not be analysed in terms of haemostatic outcome due to timings of blood sampling
Smith MP et al. 2001[18]	Bolus dose followed by continuous infusion of rFVIIa	FVII:C	Laboratory assessment	<ul style="list-style-type: none"> Target FVII:C of 10 IU/dl was found to be insufficient to prevent bleeding

FVII:C: Factor VII coagulation activity; FVIIa:C: Factor VIIa coagulation activity; FVIIa: Factor VIIa; NR: Not reported; rFVIIa: Recombinant factor VIIa

Table 5: TEG/ROTEM Analysis

Study	Haemostatic treatment	Reported factors monitored	Method of monitoring	Monitoring results
Furukawa S et al. 2015[20]	<ul style="list-style-type: none">• rFVIIa, or• aPCC	Coagulation process: <ul style="list-style-type: none">• Clotting time• Clot formation time	ROTEM	<ul style="list-style-type: none">• Clotting time and clot formation time ROTEM parameters shortened significantly after infusion of bypassing products• Clot formation time was shorter than normal in most cases after treatment with rFVIIa
Holmström M et al. 2012[22]	Bolus aPCC in combination with TXA	Whole blood coagulation profiles	ROTEM	<ul style="list-style-type: none">• During surgery, TEG showed significant improvement in CT, MaxVel and tMaxVel after aPCC and TXA and MCF increased towards normal• No significant difference in CT or MaxVel between different TXA concentrations• Significant increase in clot stability, shown by MCF, in a dose-dependent manner
Serban M et al. 2014[21]	Bolus doses and continuous infusion of: <ul style="list-style-type: none">• FVIII/FIX concentrates• rFVIIA	FVIII/FIX activity	TEG	NR

aPCC: Activated prothrombin complex concentrate; CT: Clotting time; FIX: Factor IX; FVIII: Factor VIII; MaxVel: Maximum velocity of clot formation; MCF: Maximum clot formation; NR: Not reported; rFVIIa: Recombinant factor VIIa; ROTEM: Rotational thromboelastometry; TEG: Thromboelastography; tMaxVel: Time until maximum velocity; TXA: Tranexamic acid

Table 6: Factor VIII:C

Study	Haemostatic treatment	Reported factors monitored	Method of monitoring	Monitoring results
Porcine FVIII:C				
Gatti L et al. 1984[13]	<ul style="list-style-type: none">• Bolus porcine FVIII (Hyate:C) (minor dental), or• Continuous porcine FVIII (Hyate:C) (major)	<ul style="list-style-type: none">• The antibody cross-reactivity with porcine FVIII• The relationship between preinfusion antibody titre, FVIII dosage given and its postinfusion plasma levels• The problems of 'resistance'	<ul style="list-style-type: none">• Platelet counts and haematocrits were measured with standard methods• FVIII coagulant activity measured by a one-stage method• Anti-human FVIII antibody measured in fresh plasma by the Bethesda assay method• Anti-porcine FVIII antibody measured using method based on same principles as Bethesda assay	<ul style="list-style-type: none">• Haemostatic efficacy was dependent on achieving and maintaining target levels of FVIII:C (40-50 U/dl for dental surgery)• FVIII:C was used to identify cases of 'resistance' to bypassing therapy, with treatment adjusted as appropriate
Human FVIII:C				
Danielson H et al. 2017[24]	<ul style="list-style-type: none">• Cryoprecipitate, or• Coagulation FVIII (pdFVIII or rFVIII), or• aPCC, or• rFVIIa (post-treatment switch in some individual cases)	<ul style="list-style-type: none">• FVIII:C• Development of disseminated intravascular coagulation, anaemia, or thrombocytopenia	Routine blood coagulation test	<ul style="list-style-type: none">• One patient experienced a decline in FVIII:C which led to a treatment switch
Habermann B et al. 2004[23]	Anvitoff™ (containing TXA) plus bolus FVIII or continuous rFVIIa infusion	FVIII:C levels and inhibitors	Laboratory assessment	<ul style="list-style-type: none">• A decrease of FVIII levels down to zero was measured on days 4–6 in all patients substituted with FVIII. Simultaneously an increase of the inhibitors against FVIII was noticed
Holmström M et al. 2012[22]	Bolus aPCC in combination with TXA	FVIII:C	One-stage clotting assay (FVIII activity)	NR
Serban M et al. 2014[21]	Bolus doses and continuous infusion of: <ul style="list-style-type: none">• FVIII/FIX concentrates• rFVIIa	FVIII/FIX activity	Laboratory assessment	NR

aPCC: Activated prothrombin complex concentrate; FIX: Factor IX; FVIII: Factor VIII; FVIII:C: Factor VIII coagulation activity; NR: Not reported; pdFVIII: Plasma-derived factor VIII; rFVIIa: Recombinant factor VIIa; rFVIII: Recombinant factor VIII; TXA: Tranexamic acid

Table 7: Thrombin Generation

Study	Haemostatic treatment	Factors monitored	Method of monitoring	Monitoring results
Holmström M et al. 2012[22]	Bolus aPCC in combination with TXA	LT, ETP, peak and ttPeak	TGA (on platelet-poor plasma)	<ul style="list-style-type: none">TGA showed shortened LT, ttPeak and a higher ETP and peak after aPCC + TXA administration compared to baseline, but not exceeding the values of healthy controls
Mancuso ME et al. 2016[25]	Bolus doses of: <ul style="list-style-type: none">rFVIIaaPCCSequential therapy with rFVIIa and aPCC	Platelet count	TGA	<ul style="list-style-type: none">No significant difference was found in TGA values (PRP and PPP) measured during the postoperative period by comparing procedures with (n=7) and without (n=4) bleeding complications (data not shown)

aPCC: Activated prothrombin complex concentrate; ETP: Endogenous thrombin potential; LT: Lagtime; PRP: Platelet-rich plasma; PPP: Platelet-poor plasma; rFVIIa: Recombinant factor VIIa; TGA: Thrombin generation assay; ttPeak: Time to peak; TXA: Tranexamic acid

1 **SUPPLEMENTARY DATA**

2 **Supplementary Table 1. Search Terms for MEDLINE and Embase (Searched**

3 **via the Ovid SP Platform)**

Term group	#	Searches	Hits
Patients with haemophilia	1	exp *HEMOPHILIA A/ or exp *HEMOPHILIA B/	32,171
	2	(hemophilia* or haemophilia*).tw.	49,984
	3	1 or 2	54,867
Intervention	4	(surger* or surgic* or operat* or procedure* or dental).tw.	6,877,475
	5	exp *surgical procedures, operative/	3,914,616
	6	4 or 5	8,936,274
Patients with inhibitors	7	(inhibitor* or alloantibod* or immune toler* or autoantibod*).tw.	2,648,102
Exclusion terms	8	(conference abstract or conference review).pt.	2,946,970
	9	limit 8 to yr="1946-2014"	1,907,607
	10	exp Animals/ not exp humans/	9,104,800
Combined	11	9 or 10	10,832,433
	12	3 and 6 and 7	2,925
	13	12 not 11	2,200
Total	14	remove duplicates from 13	1,383

4 MEDLINE, MEDLINE In-Process, MEDLINE Daily and MEDLINE Epub Ahead of

5 Print (1946 to present); Embase (1974 to 23 March 2018).

6

1 **Supplementary Table 2. Search Terms for The Cochrane Library (Searched**
2 **via the Wiley Online Platform)**

Term group	#	Searches	Hits
Patients with haemophilia	1	[mh "hemophilia A"] or [mh "hemophilia B"]	340
	2	(hemophilia* or haemophilia*):ti,ab,kw	1,019
	3	#1 or #2	1,019
Intervention	4	(surgery* or surgic* or operat* or procedure* or dental):ti,ab,kw	286,413
	5	[mh "surgical procedures, operative"]	122,148
	6	#4 or #5	334,058
Patients with inhibitors	7	(inhibitor* or alloantibod* or immune toler* or autoantibod*):ti,ab,kw	76,222
Total	8	#3 and #6 and #7 in Cochrane Reviews (Reviews and Protocols), Other Reviews and Trials	98

3 The Cochrane Database of Systematic Reviews (CDSR): Issue 3 of 12, March
4 2018; The Database of Abstracts of Reviews of Effects (DARE): Issue 2 of 4,
5 April 2015; The Cochrane Central Register of Controlled Trials (CENTRAL):
6 Issue 2 of 12, February 2018.

7

1 Supplementary Table 3. Search Terms for Congress Proceedings

Congress	Link	Search strategy	Hits
American Society of Hematology (ASH) Annual Meeting (2015, 2016, 2017)	2015: http://www.bloodjournal.org/content/126/23 2016: http://www.bloodjournal.org/content/128/22 2017: http://www.bloodjournal.org/content/130/Suppl_1	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Hemophil • Haemophil 	2015: 34 2016: 26 2017: 10
British Society for Haematology (BSH) Annual Scientific Meeting (2016, 2017)	2016: Abstract book was in PDF form 2017: Abstract book was in PDF form	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Hemophil • Haemophil 	2016: 6 2017: 7
European Association for Haemophilia and Allied Disorders (EAHAD) Annual Congress (2016, 2017)	2016: Abstract book was in PDF form 2017: Abstract book was in PDF form	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Surg • Operat • Proced • Dental 	2016: 54 2017: 53
Haemophilia and thrombosis research society (HTRS) Scientific Symposium (2015, 2017)	2015: Abstract book was in PDF form 2017: Abstract book was in PDF form	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Surg • Operat • Proced • Dental 	2015: 14 2017: 20
European Haemophilia Consortium (EHC) Annual Conference (2015, 2016)	<i>Abstract books for 2015 and 2016 were unavailable so were not searched</i>		
European Hematology Association (EHA) Congress (2016, 2017)	2016: Abstract book was in PDF form 2017: Abstract book was in PDF form	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Hemophil • Haemophil 	2016: 27 2017: 13
International Society for Thrombosis and Haemostasis (ISTH) Congress (2016, 2017)	2016: Abstract book was in PDF form 2017: Abstract book was in PDF form and the website was also searched (http://www.professionalabstracts.com/isth2017/iplanner/#/grid)	The following terms were searched one by one for both years: <ul style="list-style-type: none"> • Surg • Operat • Proced • Dental 	2016: 59 2017: 100
World Federation of Hemophilia (WFH) World Congress (2016)	Abstract book was in PDF form	The following terms were searched one by one: <ul style="list-style-type: none"> • Surg • Operat • Proced • Dental 	79

1 **Supplementary Table 4. Eligibility Criteria for the Systematic Review**

PICOS domain	Inclusion criteria	Exclusion criteria
Population	Haemophilia A patients with inhibitors at time of surgery Haemophilia B patients with inhibitors at time of surgery Patients with acquired haemophilia	Haemophilia A patients without inhibitors (at time of surgery) Haemophilia B patients without inhibitors (at time of surgery)
Intervention(s)	Patients undergoing surgery (haemophilia-related or unrelated procedures, including dental procedures)	Patients not undergoing surgical procedures
Comparator(s)	Any or none	No exclusion criteria
Outcomes	<p>Details of perioperative management employed in the population of interest, including:</p> <ul style="list-style-type: none"> • Monitoring of haemoglobin • Monitoring of haemostatic efficacy • Need to change dosing of haemostatic treatment or need for change in treatment • Use of thromboprophylaxis • Use of antifibrinolytics • Laboratory monitoring of global haemostasis (ROTEM, TEG, thrombin generation assessment) • Duration of treatment <p>Outcomes relating to the success of perioperative management, including:</p> <ul style="list-style-type: none"> • Bleeding control • Wound healing outcomes • Survival • Re-operation/re-admission • Infection rates • Other management-related complications 	Studies not reporting outcomes related to monitoring or management of haemophilia patients for surgery
Study design	All study designs Any study presenting original data was eligible for inclusion	Studies not presenting original data were excluded
Other considerations	Studies with abstracts or full-texts in the English language Only studies with human participants were included	Studies not published in the English language Animal studies were excluded

2 ROTEM: Rotational thromboelastometry; TEG: Thromboelastography

1 Supplementary Table 5. Studies Excluded After Full Text Screening

Study	Reason for Exclusion
Al-Salama ZT, Scott LJ. Lonoctocog Alfa: A Review in Haemophilia A. <i>Drugs</i> 2017;77:1677-1686.	Study design
Arkin S, Cooper HA, Hutter JJ, et al. Activated recombinant human coagulation factor VII therapy for intracranial hemorrhage in patients with hemophilia A or B with inhibitors: Results of the novoseven emergency-use program. <i>Haemostasis</i> 1998;28:93-98.	Wrong population
Balta A, Tornemo M, Radulovic V, et al. Monitoring of treatment with bypassing agents in patients with acquired and congenital haemophilia with inhibitors using ROTEM: A single-centre experience. <i>Haematologica</i> 2015;100:669-670.	Wrong population
Bayram I, Erbey F, Erdem S, et al. Recombinant factor VIIa and activated prothrombin-complex concentrate administration in the management of bleeding, coagulopathy and intractable coagulopathy in pediatric patients undergoing invasive medical procedures or surgery. <i>UHOD - Uluslararası Hematoloji-Onkoloji Dergisi</i> 2009;19:205-212.	Wrong population
Bedoya M, Acord M, Srinivasan A, et al. Implantable venous access devices in boys with severe hemophilia: At a tertiary pediatric institution. <i>Pediatric Radiology</i> 2017;47:S86.	Irrelevant intervention
Berger K, Frey L, Spannagl M, et al. Health economic aspects of the use of blood and blood products. [German]. <i>Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz</i> 2006;49:64-72.	Study design
Berntorp E, Astermark J, Baghaei F, et al. Treatment of haemophilia A and B and von Willebrand's disease: summary and conclusions of a systematic review as part of a Swedish health-technology assessment. <i>Haemophilia</i> 2012;18:158-65.	Study design
Biron-Andreani C, de Moerloose P, D'Oiron R, et al. Cancer detection and management in patients with haemophilia: A retrospective European multicentre study. <i>Haemophilia</i> 2013;20:78-82.	Irrelevant intervention
Birschmann I, Klamroth R, Eichler H, et al. Results of the WIRK prospective, non-interventional observational study of recombinant activated factor VII (rFVIIa) in patients with congenital haemophilia with inhibitors and other bleeding disorders. <i>Haemophilia</i> 2013;19:679-685.	Irrelevant intervention
Blanchette VS, al-Musa A, Stain AM, et al. Central venous access catheters in children with haemophilia. <i>Blood Coagulation & Fibrinolysis</i> 1996;7 Suppl 1:S39-44.	Irrelevant intervention
Boardman KP, English P. Fractures and dislocations in hemophilia. <i>Clinical Orthopaedics and Related Research</i> 1980;No. 148:221-232.	Irrelevant intervention
Bona RD, Weinstein RA, Weisman SJ, et al. The use of continuous infusion of factor concentrates in the treatment of hemophilia. <i>American Journal of Hematology</i> 1989;32:8-13.	Irrelevant intervention
Borhany M, Abid M, Fatima N, et al. Hemophilia care in Pakistan. <i>Blood Transfusion</i> 2017;15 (Supplement 3):s493.	Irrelevant intervention
Bulik O, Bulikova A, Smejkal P, et al. Preparation of patients with haemostasis disorder for dental surgery. [Czech]. <i>Vnitřní Lekarství</i> 2008;54:415-420.	Study design
Caviglia H, Candela M, Landro ME, et al. Haemophilia pseudotumours in patients with inhibitors. <i>Haemophilia</i> 2015;21:681-685.	Study design
Colvin BT. Role of plasma-exchange in the management of patients with factor VIII inhibitors. <i>La Ricerca in clinica e in laboratorio</i> 1983;13:85-93.	Study design
Cooper HA, Gilchrist GS, Hoots WK, et al. Comparison of two doses of recombinant factor VIIa (rFVIIa) for producing hemostasis during and after surgery in patients (PTS) with hemophilia A or B and inhibitors and PTS with acquired inhibitors. <i>Blood</i> 1997;90:600a.	Study design

Study	Reason for Exclusion
Coppola A, Minno M, Tufano A, et al. Treatment for preventing bleeding in people with congenital bleeding disorders undergoing surgery: A systematic review of randomised controlled trials. <i>Thrombosis research</i> . Volume 134, 2014:S4-s5.	Study design
Coppola A, Windyga J, Tufano A, et al. Treatment for preventing bleeding in people with haemophilia or other congenital bleeding disorders undergoing surgery. <i>The Cochrane database of systematic reviews</i> 2015;2:CD009961.	Study design
Dargaud Y, Lienhart A, Negrier C. Prospective assessment of thrombin generation test for dose monitoring of bypassing therapy in hemophilia patients with inhibitors undergoing elective surgery. <i>Blood</i> 2010;116:5734-5737.	Study design
Dargaud Y, Pavlova A, Lacroix-Desmazes S, et al. Achievements, challenges and unmet needs for haemophilia patients with inhibitors. <i>Haemophilia</i> 2016;22:1-24.	Study design
Dekoven M, Wisniewski T, Petrilla A, et al. Patient/caregiver perceived benefits and barriers to elective orthopedic surgery (EOS) in patients with congenital hemophilia with inhibitors. <i>Journal of Medical Economics</i> 2012;15:305-312.	Irrelevant intervention
Domm JA, Hudson MG, Janco RL. Complications of central venous access devices in paediatric haemophilia patients. <i>Haemophilia</i> 2003;9:50-56.	Irrelevant intervention
Economou M, Teli A, Adremerina A, et al. Absence of thrombotic complications with the use of bypassing agents in young hemophilia patients with inhibitor presence. <i>Haemophilia</i> 2018;24 (Supplement 1):105.	Irrelevant intervention
Escobar M, Maahs J, Hellman E, et al. Multidisciplinary management of patients with haemophilia with inhibitors undergoing surgery in the United States: Perspectives and best practices derived from experienced treatment centres. <i>Haemophilia</i> 2012;18:971-981.	Study design
Furukawa S, Nogami K, Ogiwara K, et al. Systematic monitoring of hemostatic management in hemophilia A patients with inhibitor in the perioperative period using rotational thromboelastometry. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:350.	Duplicate
Galstian GM, Spirin M, Zozulya N, et al. Providing hemostasis for long-term central venous access device (LTCVAD) placement in patients with factor VIII (FVIII) inhibitors. <i>Blood</i> . Conference: 59th Annual Meeting of the American Society of Hematology, ASH 2017;130.	Irrelevant intervention
Ghosh K, Shetty S, Jijina F, et al. Role of epsilon amino caproic acid in the management of haemophilic patients with inhibitors. <i>Haemophilia</i> 2004;10:58-62.	Wrong population
Givol N, Hirschhorn A, Lubetsky A, et al. Oral surgery-associated postoperative bleeding in haemophilia patients - a tertiary centre's two decade experience. <i>Haemophilia</i> 2015;21:234-240.	Irrelevant intervention
Goodnight Jr SH, Common HH, Lovrien EW. Factor VIII inhibitor following surgery for epidural hemorrhage in hemophilia: successful therapy with a concentrate containing factors II, VII, IX, and X. <i>Journal of Pediatrics</i> 1976;88:357-358.	Wrong population
Gozden HE, Ozkalemkas F, Ozkocaman V, et al. Evaluation of patients with hemophilia; Uludag University experience. <i>Thrombosis Research</i> 2016;141:S37.	Wrong population
Haque Q, Feng X, Abuduaini Y. Intracranial haemorrhage in children with inherited bleeding disorders: A single center study in China. <i>Hong Kong Journal of Paediatrics</i> 2018;23(1):69.	Irrelevant intervention
Haque Q, Li C, Abuduaini Y, et al. Intracranial hemorrhage in children with inherited bleeding disorders: A single center study in China. <i>Blood</i> . Conference: 59th Annual Meeting of the American Society of Hematology, ASH 2017;130.	Duplicate
Hay CRM, Negrier C, Ludlam CA. The treatment of bleeding in acquired haemophilia with recombinant factor VIIa: A multicentre study. <i>Thrombosis and Haemostasis</i> 1997;78:1463-1467.	Irrelevant intervention
Hedner U. Factor VIIa in the treatment of haemophilia. <i>Blood coagulation & fibrinolysis: an international journal in haemostasis and thrombosis</i> 1990;1:307-317.	Study design

Study	Reason for Exclusion
Hirose J, Takedani H, Koibuchi T. The risk of elective orthopaedic surgery for haemophilia patients: Japanese single-centre experience. <i>Haemophilia</i> 2013;19:951-955.	Irrelevant intervention
Holmstrom M, Astermark J, Brodin E, et al. Swedish national registry for bleeding disorders-first report. <i>Haemophilia</i> 2018;24 (Supplement 1):101.	Wrong population
Hvid I, Rodriguez-Merchan EC. Orthopaedic surgery in haemophilic patients with inhibitors: An overview. <i>Haemophilia</i> 2002;8:288-291.	Study design
Jones ML, Wight J, Paisley S, et al. Control of bleeding in patients with haemophilia A with inhibitors: A systematic review. <i>Haemophilia</i> 2003;9:464-520.	Study design
Kenet G, Lubetsky A, Gitel S, et al. Treatment of bleeding episodes in patients with hemophilia and an inhibitor: Comparison of two treatment protocols with recombinant activated factor VII. <i>Blood Coagulation and Fibrinolysis</i> 2000;11:S35-S38.	Wrong population
Kleinschmidt S, Plinkert PK, Fuchs-Buder T, et al. [Haemostatic disorders in ENT patients.Part 2: Pathophysiology, diagnostics, clinical feature and therapy]. <i>HNO</i> 2003;51:251-266.	Study design
Klintman J, Berntorp E. Epidemiological aspects of inhibitor development in hemophilia and strategies of management. <i>Expert Opinion on Orphan Drugs</i> 2016;4:153-168.	Study design
Klukowska A, Laguna P, Rawicz M. Procedures for CV catheters insertion in children with congenital coagulation disorders. [Polish]. <i>Medycyna wieku rozwojowego</i> 2008;12:1126-1129.	Irrelevant intervention
Kreuz W, Gill JC, Rothschild C, et al. Full-length sucrose-formulated recombinant factor VIII for treatment of previously untreated or minimally treated young children with severe haemophilia A: Results of an international clinical investigation. <i>Thrombosis and Haemostasis</i> 2005;93:457-467.	Wrong population
Kulkarni R, Presley RJ, Lusher JM, et al. Complications of haemophilia in babies (first two years of life): a report from the Centers for Disease Control and Prevention Universal Data Collection System. <i>Haemophilia</i> 2017;23:207-214.	Wrong population
Laguna P, Klukowska A. Management of oral bleedings with recombinant factor VIIa in children with haemophilia A and inhibitor. <i>Haemophilia</i> 2005;11:2-4.	Study design
Liesner RJ, Abashidze M, Alenikova O, et al. Immunogenicity, efficacy and safety of Nuwiq ^{<sup></sup>} (human-cl rhFVIII) in previously untreated patients with severe haemophilia A-Interim results from the NuProtect Study. <i>Haemophilia</i> 2017;16:16.	Irrelevant intervention
Lim MY, Nielsen B, Ma A, et al. Clinical features and management of haemophilic pseudotumours: A single US centre experience over a 30-year period. <i>Haemophilia</i> 2013;20:e58-e62.	Study design
Lulla RR, Allen GA, Zakarija A, et al. Transplacental transfer of postpartum inhibitors to factor VIII. <i>Haemophilia</i> 2010;16:14-17.	Wrong population
Lusher JM, Lee CA, Kessler CM, et al. The safety and efficacy of B-domain deleted recombinant factor VIII concentrate in patients with severe haemophilia A. <i>Haemophilia</i> 2003;9:38-49.	Irrelevant intervention
Makris M. Systematic review of the management of patients with haemophilia A and inhibitors. <i>Blood Coagulation and Fibrinolysis</i> 2004;15:S25-S27.	Study design
McPherson J, Teague L, Lloyd J, et al. Experience with recombinant factor VIIa in Australia and New Zealand. <i>Haemostasis</i> 1996;26:109-117.	Study design
Mingot-Castellano ME, Perez-Montes R, Canaro M, et al. Successful treatment of bleeding in acquired hemophilia A with activated prothrombin complex concentrate in Spain. <i>Blood. Conference: 59th Annual Meeting of the American Society of Hematology, ASH</i> 2017;130.	Irrelevant intervention

Study	Reason for Exclusion
Mortazavi SMJ, Najafi A, Toogeh G. Total joint replacement in haemophilia A patients with high titre of inhibitor using a new brand recombinant factor VIIa (Aryoseven). Haemophilia 2016;22:e451-e453.	Study design
Negrier C, Ragni MV, Georgiev P, et al. Perioperative management in patients with hemophilia receiving fitusiran, an investigational rna therapeutic targeting antithrombin for the treatment of hemophilia. Blood. Conference: 59th Annual Meeting of the American Society of Hematology, ASH 2017;130.	Duplicate
Nilsson IM, Hedner U. Immunosuppressive treatment in haemophiliacs with inhibitors to factor VIII and factor IX. Scandinavian Journal of Haematology 1976;16:369-382.	Study design
Obergfell A, Auvinen MK, Mathew P. Recombinant activated factor VII for haemophilia patients with inhibitors undergoing orthopaedic surgery: A review of the literature. Haemophilia 2008;14:233-241.	Study design
Parameswaran R, Shapiro AD, Gill JC, et al. Dose effect and efficacy of rFVIIa in the treatment of haemophilia patients with inhibitors: Analysis from the Hemophilia and Thrombosis Research Society Registry. Haemophilia 2005;11:100-106.	Irrelevant intervention
Park YS. Preferences between surgical or medical treatment in hemophilia patients with spontaneous epidural hematoma may vary. Haemophilia 2018;24 (Supplement 1):133.	Irrelevant intervention
Pruthi RK, Mathew P, Valentino LA, et al. An open-label, randomized, parallel, multi-center trial comparing the safety and efficacy of rFVIIa when administered as IV bolus or IV continuous infusion to hemophilia patients with inhibitors during and after surgery. Blood 2004;104:3975.	Irrelevant intervention
Rodriguez-Merchan EC. Surgery in haemophilic patients with inhibitors. Haemophilia 2004;10 Suppl 2:1-2.	Duplicate
Rodriguez-Merchan EC, Wiedel JD, Wallny T, et al. Elective orthopaedic surgery for inhibitor patients. Haemophilia 2003;9:625-31.	Duplicate
Rudowski WJ. Major surgery in hemophilia. Annals of the Royal College of Surgeons of England 1981;63:111-117.	Irrelevant intervention
Salaj P, Louzil J, Geierova V, et al. Diagnosis and management of acquired haemophilia-single centre experience. Journal of Thrombosis and Haemostasis 2016;14:56.	Irrelevant intervention
Salcioglu Z, Sen HS, Tugcu D, et al. Congenital factor deficiencies: Twenty-five-year follow-up. Journal of Thrombosis and Haemostasis 2015;13:358-359.	Irrelevant intervention
Salzmann G, Schramm W, Feifel G. The hemophiliac as a surgical patient. [German]. Munchener Medizinische Wochenschrift 1977;119:677-684.	Wrong population
Schoppmann A, Jaeger K, Berg R, et al. Review of the literature of FEIBA administration in patients with hemophilia B and inhibitors. Journal of Coagulation Disorders 2011;3:14-26.	Study design
Schulz, S. Inhibitor hemophilia in oral surgery. [German]. Zahn-, Mund-, und Kieferheilkunde mit Zentralblatt 1984;72:824-848.	Non-English language
Schwartz RS, Abildgaard CF, Aledort LM, et al. Human recombinant DNA-derived antihemophilic factor (factor VIII) in the treatment of hemophilia A. New England Journal of Medicine 1990;323:1800-1805.	Wrong population
Seaman CD, Ragni MV. Sequential bypassing agents during major orthopedic surgery: a new approach to hemostasis. Blood Advances 2017;1:1309-1311.	Study design
See A, Sudirman SR, Huang XY. Spontaneous multilevel airway haemorrhage in acquired haemophilia A. European Archives of Oto-Rhino-Laryngology 2016:1-4.	Study design
Serban M, Mihailov D, Pop L, et al. Development of inhibitors in haemophilia. Hamostaseologie 2011;31:S20-S23.	Wrong population

Study	Reason for Exclusion
Serban M, Ursu E, Cernat L, et al. Thrombin generation and whole blood viscoelastic assays in the monitoring of haemophilia with inhibitors. <i>Haematologica</i> 2016;101:407.	Irrelevant intervention
Sholzberg M, Phua C, Tsui H, et al. Heparin and protamine confound factor activity and inhibitor testing while on cardiopulmonary bypass. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:807.	Wrong population
Shutov SA, Kovalenko AV, Soboleva OA, et al. [Surgical treatment of the complication of urolithiasis in patients with inhibitor form of hemophilia]. <i>Khirurgiia</i> 2017:104-107.	Non-English language
Smith MP, Giangrande P, Pollman H, et al. A postmarketing surveillance study of the safety and efficacy of ReFacto (St Louis-derived active substance) in patients with haemophilia A. <i>Haemophilia</i> 2005;11:444-451.	Wrong population
Stachnik JM, Gabay MP. Continuous infusion of coagulation factor products. <i>Annals of Pharmacotherapy</i> 2002;36:882-891.	Study design
Stine KC, Shrum D, Becton DL. Use of FEIBA for invasive or surgical procedures in patients with severe hemophilia A or B with inhibitors. <i>Journal of Pediatric Hematology/Oncology</i> 2007;29:216-221.	Study design
Tjonnfjord GE, Brinch L, Gedde-Dahl III, et al. Activated prothrombin complex concentrate (FEIBA) treatment during surgery in patients with inhibitors to FVIII/IX. <i>Haemophilia</i> 2004;10:174-178.	Duplicate
Tuinenburg A, Damen SAJ, Ypma PF, et al. Cardiac catheterization and intervention in haemophilia patients: Prospective evaluation of the 2009 institutional guideline. <i>Haemophilia</i> 2013;19:370-377.	Study design
Valentino LA, Cooper DL, Goldstein B. Surgical Experience with rFVIIa (NovoSeven) in congenital haemophilia A and B patients with inhibitors to factors VIII or IX. <i>Haemophilia</i> 2011;17:579-589.	Study design
van Veen JJ, Maclean RM, Hampton KK, et al. Major surgery in severe haemophilia A with inhibitors using a recombinant factor VIIa and activated prothrombin complex concentrate hybrid regimen. <i>Haemophilia</i> 2014;20:587-592.	Study design
Varon D, Martinowitz U. Continuous infusion therapy in haemophilia. <i>Haemophilia</i> 1998;4:431-435.	Study design
Watts RG, Cook RP. Operative management and outcomes in children with congenital bleeding disorders: A retrospective review at a single haemophilia treatment centre. <i>Haemophilia</i> 2012;18:421-425.	Irrelevant intervention
Wensley RT, Stevens RF, Burn AM, et al. Plasma exchange and human factor VIII concentrate in managing haemophilia A with factor VIII inhibitors. <i>British Medical Journal</i> 1980;281:1388-1389.	Wrong population
White GC, 2nd. Seventeen years' experience with Autoplex/Autoplex T: evaluation of inpatients with severe haemophilia A and factor VIII inhibitors at a major haemophilia centre. <i>Haemophilia</i> 2000;6:508-12.	Irrelevant intervention
Windyga J, Stefanska-Windyga E, Odnoczko E, et al. Activated prothrombin complex concentrate in combination with tranexamic acid: a single centre experience for the treatment of mucosal bleeding and dental extraction in haemophilia patients with inhibitors. <i>Haemophilia</i> 2016;22:e465-e468.	Study design
Young G, Sidonio RF, Liesner R, et al. HAVEN 2 updated analysis: Multicenter, open-label, phase 3 study to evaluate efficacy, safety and pharmacokinetics of subcutaneous administration of emicizumab prophylaxis in pediatric patients with hemophilia a with inhibitors. <i>Blood. Conference: 59th Annual Meeting of the American Society of Hematology, ASH</i> 2017;130.	Irrelevant intervention

1 **Supplementary Table 6. Case Studies Excluded After Abstract Screening**

Study
Abajas YL, Monahan PE, Neufeld EJ, et al. Novel approach for immunosuppression and peri-operative recombinant porcine factor viii replacement for pediatric congenital hemophilia a with high-titer inhibitor. American Journal of Hematology 2016;91 (9):E416.
Abildgaard CF, Penner JA, Watson-Williams EJ. Anti-inhibitor coagulant complex (Autoplex) for treatment of factor VIII inhibitors in hemophilia. Blood 1980;56:978-984.
Al-Trabolsi HA. Factor VIIa overdose: Clinical and laboratory observations. Current Pediatric Research 2004;8:19-21.
Ambaglio C, Preti PS, Sacco C, et al. Orthopaedic surgery in a patient affected by severe haemophilia a with inhibitors: Three surgeries-three different responses to bypassing agents. Haemophilia 2017;23:112-113.
Amici JM. Surgery of basal cell carcinoma in a man with hemophilia: Full thickness skin graft. [French]. Nouvelles Dermatologiques 2004;23:130-131.
Antic D, Elezovic I, Sufajdzic N, et al. Application of recombinant activated factor VII in treatment of intracranial haemorrhage in haemophilic patient with inhibitor. [Croatian]. Srpski arhiv za celokupno lekarstvo 2008;136 Suppl 3:218-221.
Aouba A, Dezamis E, Sermet A, et al. Uncomplicated neurosurgical resection of a malignant glioneuronal tumour under haemostatic cover of rFVIIa in a severe haemophilia patient with a high-titre inhibitor: A case report and literature review of rFVIIa use in major surgeries. Haemophilia 2010;16:54-60.
Apter B, McCarthy V, Shapiro SS, et al. Successful preoperative apheresis of factor VIII antibody using factor VIII concentrate as a replacement fluid. Journal of Clinical Apheresis 1986;3:140.
Ashrani AA, Reding MT, Shet A, et al. Successful liver transplantation in a patient with severe haemophilia A and a high-titre factor VIII inhibitor. Haemophilia 2004;10:735-737.
Banov L, Pavanello M, Piattelli G, et al. Successful urgent neurosurgery management with rFVIIa mega doses in a child with haemophilia A and high titre inhibitor. Blood Coagulation and Fibrinolysis 2014;25:518-521.
Barbara DW, McKenzie KM, Parikh SA, et al. Successful perioperative management of severe bleeding from undiagnosed acquired factor VIII inhibitors. Journal of Cardiothoracic and Vascular Anesthesia 2015;29:731-734.
Batorova A, Morongova A, Tagariello G, et al. Challenges in the management of hemophilia B with inhibitor. Seminars in Thrombosis and Hemostasis 2013;39:767-771.
Bell BA, Birch K, Glazer S. Experience with recombinant factor VIIA in an infant hemophiliac with inhibitors to FVIII:C undergoing emergency central line placement: A case report. American Journal of Pediatric Hematology/Oncology 1993;15:77-79.
Bennetts NA, Mergelmeyer JE, Reimer EJ, et al. Initial Manifestation of Acquired Hemophilia A After a Routine Tooth Extraction. A Case Report and Literature Review. Journal of Oral and Maxillofacial Surgery 2018;76:490-494.
Berlocher WC, King DL. Considerations in the dental management of the factor VIII-deficient child with inhibitors. Pediatric dentistry 1979;1:188-191.
Bhave A, Srivastava A, Lee V, et al. Low-dose activated factor IX complex concentrates (FEIBA(R)) for post-operative haemostasis in a patient with high responding factor VIII inhibitors. Haemophilia 1995;1:274-6.
Biron-Andreani C, Dupeyron G, Mainemer M, et al. Successful use of recombinant factor VIIa in a haemophiliac with inhibitor undergoing cataract surgery. Blood Coagulation and Fibrinolysis 2001;12:215-216.
Blatt PM, Pearsall AH, Givhan EG, et al. Haemostatic failure of prothrombin complex concentrates during elective dental procedure. Thrombosis and Haemostasis 1980;42:1604-1606.

Study
Bona RD, Pasquale DN, Kalish RI, et al. Porcine factor VIII and plasmapheresis in the management of hemophiliac patients with inhibitors. <i>American Journal of Hematology</i> 1986;21:201-207.
Bontempo FA, Lewis JH, Spero JA, et al. Heart transplant in a hemophiliac with an acquired factor VIII inhibitor: Synthesis of factor VIII:C in pericardial fluid. <i>Transplantation Proceedings</i> 1988;20:790-791.
Boughton BJ, Payne A, Serman A, et al. Elective surgery in a haemophilic patient with high titre inhibitors: use of extracorporeal protein A immunoabsorption. <i>Journal of Clinical Pathology</i> 1990;43:172.
Brackmann HH, Effenberger W, Hess L, et al. Immune tolerance induction: A role for recombinant activated factor VII (rFVIIa)? <i>European Journal of Haematology, Supplement</i> 1998;61:18-23.
Briet E, van der Meer FW, van Dijk-van Kempen CJ, et al. Sequential administration of human and porcine factor VIII for surgical treatment of a parotid tumour in a patient with a factor VIII inhibitor. <i>Acta Haematologica</i> 1985;73:97-100.
Burk CD, Miller L, Handler SD, et al. Preoperative history and coagulation screening in children undergoing tonsillectomy. <i>Pediatrics</i> 1992;89:691-695.
Byhahn C, Lischke V, Westphal K. Translaryngeal tracheostomy in highly unstable patients. <i>Anaesthesia</i> 2000;55:678-682.
Candiotto L, Fullone FW, Ricciardi A, et al. Correction of knee flexion contracture at the time of surgical fixation of a femoral supracondylar fracture in a haemophiliac with inhibitors. <i>Blood Transfusion</i> 2015;13:333-335.
Carr Jr ME, Loughran TP, Cardea JA, et al. Successful use of recombinant factor VIIa for hemostasis during total knee replacement in a severe hemophiliac with high-titer factor VIII inhibitor. <i>International journal of hematology</i> 2002;75:95-99.
Castillo-Canadas AM, Serrano-Diana C, Lopez-Del Cerro E, et al. Diagnosis and treatment of hemophilia A acquired during postpartum. [Spanish]. <i>Ginecologia y Obstetricia de Mexico</i> 2014;82:688-696.
Caviglia H, Landro ME, Galatro GA, et al. Platelet rich in fibrin (PRF) in hemophilia. <i>Haemophilia</i> 2018;24 (Supplement 1):83.
Chau A, Wu J, Ansermino M, et al. A Jehovah's Witness child with hemophilia B and factor IX inhibitors undergoing scoliosis surgery. <i>Canadian Journal of Anesthesia</i> 2008;55:47-51.
Chen YC, Chang JY, Hsueh EJ, et al. Acquired hemophilia A: Report of two cases. <i>Chinese Medical Journal (Taipei)</i> 1998;61:538-544.
Chu M, Li H. Regional anesthesia in acquired hemophilia a (factor viii inhibitor positive). <i>Regional Anesthesia and Pain Medicine</i> . Conference: 41st Annual Regional Anesthesiology and Acute Pain Medicine Meeting of the American Society of Regional Anesthesia and Pain Medicine, ASRA 2016;41.
Chuansumrit A, Hathirat P, Keorochana S, et al. Disarticulation of a knee joint in a haemophiliac with high inhibitor titre. <i>Haemophilia</i> 1996;2:116-119.
Cooper HA, Jones CP, Campion E, et al. Rationale for the use of high dose rFVIIa in a high-titre inhibitor patient with haemophilia B during major orthopaedic procedures. <i>Haemophilia</i> 2001;7:517-522.
Croteau SE, Abajas YL, Wolberg AS, et al. Recombinant porcine factor VIII for high-risk surgery in paediatric congenital haemophilia A with high-titre inhibitor. <i>Haemophilia</i> 2017;23:e93-e98.
Damodar S, Bhat P, Kumar P, et al. Successful Aortic Valve Replacement Surgery in a Patient with Severe Haemophilia a with Low Titre Inhibitor. <i>Indian Journal of Hematology and Blood Transfusion</i> 2014;30:64-66.
Dargaud Y, Lienhart A, Meunier S, et al. Major surgery in a severe haemophilia A patient with high titre inhibitor: Use of the thrombin generation test in the therapeutic decision. <i>Haemophilia</i> 2005;11:552-558.
DeWitt RT, Feinstein DI. Prothrombin complex concentrate. Use in a hemophiliac with a factor VIII inhibitor. <i>Archives of Internal Medicine</i> 1977;137:1211-1213.

Study
Di Gaetano R, Belvini D, Salviato R, et al. Flow cytometry evaluation of INF-gamma and il-10 synthesis by T lymphocytes in a haemophilia a patient with inhibitor after infusion of FVIII. <i>Blood Transfusion</i> 2017;15 (Supplement 4):s545.
Divanon F, Hecquard C, Borel-Derlon A. Experience with use of recombinant activated factor VII. <i>Journal of Clinical Pharmacy and Therapeutics</i> 2002;27:133-138.
Dolatkhah R, Bazavar MR, Poureisa M, et al. Successful management of total knee replacement in a high responder hemophilia patient with a history of inhibitor. <i>Iranian Red Crescent Medical Journal</i> 2013;15:18-20.
Doughty HA, Coles J, Parmar K, et al. The successful removal of a bleeding intracranial tumour in a severe haemophiliac using an adjusted dose continuous infusion of monoclonal factor VIII. <i>Blood Coagulation and Fibrinolysis</i> 1995;6:31-34.
Doughty HA, Northeast A, Sklair L, et al. The use of recombinant factor VIIa in a patient with acquired haemophilia A undergoing surgery. <i>Blood Coagulation and Fibrinolysis</i> 1995;6:125-128.
Durham TM, Hodges ED, Harper J, et al. Management of traumatic oral-facial injury in the hemophiliac patient with inhibitor: case report. <i>Pediatric dentistry</i> 1993;15:282-287.
Eigner TL. Use of intraligamentary anesthesia in a patient with severe hemophilia and factor VIII inhibitor. <i>Special care in dentistry : official publication of the American Association of Hospital Dentists, the Academy of Dentistry for the Handicapped, and the American Society for Geriatric Dentistry</i> 1990;10:121-124.
Evans BE, Irving SP, Aledort LM. Use of microcrystalline collagen for hemostasis after oral surgery in a hemophiliac. <i>Journal of Oral Surgery</i> 1979;37:126-8.
Faradji A, Bonnomet F, Lecocq J, et al. Knee joint arthroplasty in a patient with haemophilia A and high inhibitor titre using recombinant factor VIIa (NovoSeven): A new case report and review of the literature. <i>Haemophilia</i> 2001;7:321-326.
Feistritzer C, Wildner SM, Wurtinger P, et al. Successful immune tolerance induction using turoctocog alfa in an adult haemophilia A patient. <i>Blood Coagulation and Fibrinolysis</i> 2017;28:181-184.
Forsyth A, Zourikian N. How we treat: Considerations for physiotherapy in the patient with haemophilia and inhibitors undergoing elective orthopaedic surgery. <i>Haemophilia</i> 2012;18:550-553.
Franchini M, Capra F, Capelli C, et al. Clinical efficacy of recombinant activated factor VII (rFVIIa) during acute bleeding episode and surgery in a patient with acquired hemophilia A with high inhibitor titer. <i>Haematologica</i> 2001;86:E12.
Frauchiger LH, Harstall R, Kajahn J, et al. Bilateral total knee arthroplasty in a patient with haemophilia A, high inhibitor titre and aneurysma spurium of the popliteal artery: A case report. <i>Swiss Medical Weekly</i> 2010;140.
Gay ND, Azar SS, Salomon O, et al. Management of a patient with factor XI deficiency with inhibitors undergoing cardiac surgery: A case report and review of the literature. <i>American Journal of Hematology</i> 2016;91 (9):E413.
Gillet B, Sigaud M, Ternisien C, et al. Can you predict the clinical efficacy of bypassing agents in patients with hemophilia and current inhibitor? Discussion about a case report in cardiovascular surgery. <i>Haemophilia</i> 2018;24 (Supplement 1):58.
Giuffrida G, Lombardo R, Parrinello NL, et al. Percutaneous transluminal angioplasty and stent implantation for aortic coarctation in haemophilia A patient with high-titre factor VIII inhibitors. <i>Haemophilia</i> 2014;20:e336-e338.
Goddard N. Case studies: Orthopaedic surgery in adult patients with haemophilia A with inhibitors. <i>Haemophilia, Supplement</i> 2005;11:32-37.
Gomes H, Martinez F, Robelo B, et al. Delayed diagnosis of pulmonary embolism in a patient with a prior history of acquired haemophilia. <i>Haemophilia</i> 2015;21:39.
Gonzalez S, Mai S, Jahng A. Intractable lower GI bleeding due to acquired hemophilia a in a patient with rectal cancer. <i>American Journal of Gastroenterology</i> 2017;112 (Supplement 1):S1045-S1047.

Study
Goodknight SH, Common HH, Lovrein EW. Letter: Factor VIII inhibitor following surgery for epidural hemorrhage in hemophilia: successful therapy with a concentrate containing factors II, VII, IX, and X. <i>The Journal of pediatrics</i> 1976;88:356-357.
Gopalakrishnan N, Usha T, Thopalan B, et al. Hemodialysis in a patient with severe hemophilia A and factor VIII inhibitor. <i>Hemodialysis International</i> 2016;20:E11-E13.
Gorska-Kosicka M, Paluszkiewicz P, Krasowska D, et al. Uncontrolled postoperative bleeding in woman with pemphigus and undiagnosed acquired haemophilia A. [Polish]. <i>Hematologia</i> 2013;4:71-75.
Goudemand J, Tagariello G, Lopaciuk F. Cases of surgery in high-responder haemophilia patients. <i>Haemophilia</i> , Supplement 2004;10:46-49.
Gregg R, Lester W, Bramhall S, et al. Orthotopic liver transplantation in a patient with severe haemophilia A and a high-titre factor VIII inhibitor from an antithrombin-deficient cadaveric donor. <i>Haemophilia</i> 2013;19:e96-e97.
Griffen AL, Hoots WK, Carter AB. Activated prothrombin complex concentrates in the management of the hemophiliac with Factor VIII inhibitor: case report. <i>Pediatric Dentistry</i> 1987;9:321-4.
Gringeri A, Santagostino E, Mannucci PM. Failure of recombinant activated factor VII during surgery in a hemophiliac with high-titer factor VIII antibody. <i>Haemostasis</i> 1991;21:1-4.
Guglielmone H, Jarchum G, Minoldo S. Liver transplantation in a patient with mild haemophilia A and low-titres of factor VIII inhibitors treated with recombinant factor VIIa. The first Argentinean case. <i>Haemophilia</i> 2011;17:317-318.
Hanna WT, Madigan RR, Miles MA, et al. Activated factor IX complex in treatment of surgical cases of hemophilia A with inhibitors. <i>Thrombosis and Haemostasis</i> 1981;46:638-641.
Hasson DM, Poole AE, de la Fuente B, et al. The dental management of patients with spontaneous acquired factor VIII inhibitors. <i>Journal of the American Dental Association (1939)</i> 1986;113:633-636.
Hayashi T, Morishita E, Asakura H, et al. Two cases of acquired hemophilia A in elderly patients. [Japanese]. <i>Japanese Journal of Geriatrics</i> 2010;47:329-333.
Heiland M, Weber M, Schmelzle R. Life-Threatening Bleeding After Dental Extraction in a Hemophilia A Patient With Inhibitors to Factor VIII: A Case Report. <i>Journal of Oral and Maxillofacial Surgery</i> 2003;61:1350-1353.
Heisel MA, Gomperts ED, McComb JG, et al. Use of activated prothrombin complex concentrate over multiple surgical episodes in a hemophilic child with an inhibitor. <i>Journal of Pediatrics</i> 1983;102:951-954.
Ho LP, Ho YK, Tien SL. Case report: Induction of immune tolerance to factor VIII inhibitor after a major operation. <i>Annals of the Academy of Medicine Singapore</i> 2007;36:431-434.
Horton S, Martlew V, Wilde J, et al. Re-emergence of a low-titre factor VIII inhibitor after liver transplant. <i>Haemophilia</i> 2012;18:e69-e71.
Hutchinson RJ, Penner JA, Hensinger RN. Anti-inhibitor coagulant complex (autoplex) in hemophilia inhibitor patients undergoing synovectomy. <i>Pediatrics</i> 1983;71:631-633.
Ichikawa S, Kohata K, Okitsu Y, et al. Acquired hemophilia A with sigmoid colon cancer: Successful treatment with rituximab followed by sigmoidectomy. <i>International Journal of Hematology</i> 2009;90:33-36.
Ilg A, Stahlschmidt K, Zotz RB, et al. Interdisciplinary management of total knee replacement in a haemophilia patient with high-titre inhibitor and severe arthropathy complicated by an aneurysmatic bone cyst. <i>Haemophilia</i> 2009;15:377-9.
Ilkhchoui Y, Koshkin E, Windsor JJ, et al. Perioperative management of acquired hemophilia a: a case report and review of literature. <i>Anesthesiology & Pain Medicine</i> 2014;4:e11906.

Study
Isobe K, Koh K, Uehara T, et al. Successful sequential therapy using rfvii with a novel agent (plasma-derived factor fviia and factor fx mixture) to control perioperative bleeding in a patient with severe hemophilia a and inhibitors. <i>Haematologica</i> 2016;101:709.
Janic D, Brdar R, Krstic Z, et al. Successful concurrent triple surgery in an adolescent patient with haemophilia A and inhibitors treated with recombinant factor VIIa [2]. <i>Haemophilia</i> 2007;13:214-216.
Janic D, Brdar R, Krstic Z, et al. Successful concurrent triple surgery in an adolescent patient with haemophilia A and inhibitors treated with recombinant factor VIIa [4]. <i>Haemophilia</i> 2007;13:447-449.
Janic D, Petronic I, Dokmanovic L, et al. Rehabilitation in haemophilic children with inhibitors using recombinant activated factor VII. [Croatian]. <i>Srpski arhiv za celokupno lekarstvo</i> 2008;136 Suppl 3:226-230.
Jentsch T, Brand-Staufer B, Schafer FP, et al. Illustrated operative management of spontaneous bleeding and compartment syndrome of the lower extremity in a patient with acquired hemophilia A: A case report. <i>Journal of Medical Case Reports</i> 2014;8 (1) (no pagination).
Jones AE, Roy A, Armstrong T, et al. Successful liver surgery in a haemophilia patient with high titre factor VIII inhibitor. <i>Haemophilia</i> 2009;15:1332-1333.
Kanyike FB, Abdul-Salam SA, Prakash B, et al. Use of recombinant factor VIIa (NovoSeven) in a haemophilia A patient with inhibitor in Kuwait. <i>Haemophilia</i> 1999;5:273-275.
Kashyap R, Choudhry VP, Mahapatra M, et al. Postpartum acquired haemophilia: Clinical recognition and management. <i>Haemophilia</i> 2001;7:327-330.
Kawasaki Y, Saeki N, Kawamoto M, et al. Perioperative use of recombinant activated factor VII (rF VIIA) in a patient with hemophilia A having inhibitors. [Japanese]. <i>Japanese Journal of Anesthesiology</i> 2005;54:926-928.
Kaya Z, Orhan O, Turanli S, et al. Successful total hip replacement with sequential administration of bypassing agents in an adolescent boy with hemophilia A and high inhibitor titers. <i>Blood Coagulation & Fibrinolysis</i> 2017;12:12.
Kaya Z, Orhan O, Turanli S, et al. Total HIP replacement in a hemophilia-a patient with high titer inhibitor. <i>Blood</i> 2015;126 (23):4707.
Keller A, Terrier F, Schneider PA, et al. Pelvic haemophilic pseudotumour: Management of a patient with high level of inhibitors. <i>Skeletal Radiology</i> 2002;31:550-553.
Khakhar AK, Chan NG, Allan DS, et al. Catastrophic microangiopathy induced by high-titre factor VIII inhibitors after liver transplantation for haemophilia A with cirrhosis. <i>Haemophilia</i> 2005;11:623-628.
Klukowska A, Laguna P, Obitko-Pludowska A, et al. Use of activated recombinant factor VII (rFVIIa) in the treatment of hemophilia A children with high-titer inhibitor. [Polish]. <i>Acta Haematologica Polonica</i> 1998;29:401-406.
Konkle BA, Nelson C, Forsyth A, et al. Approaches to successful total knee arthroplasty in haemophilia A patients with inhibitors. <i>Haemophilia</i> 2002;8:706-10.
Kovalova Z, Shelkova S, Cebura E, et al. Unconvincing reduction of inhibitor titers during first Latvian ITI experience with high-dosage coagulation factor regime. <i>Haemophilia</i> 2018;24 (Supplement 1):135.
Koyama T, Nagao T, Tsunozaki H, et al. Successful management of massive intraperitoneal bleeding in a hemophilia A patient with inhibitor by surgical debridement of the incomplete hematoma and administration of recombinant factor VIII and activated factor VII. <i>Pathophysiology of Haemostasis and Thrombosis</i> 2006;35:405-407.
Kubisz P, Pjamenova I, Stasko J, et al. Use of secondary prophylaxis with recombinant activated factor VII in haemophilia a with inhibitor: Our experience and the review of the literature. [Slovak]. <i>Transfuzie a Hematologie Dnes</i> 2009;15:210-215.
Lauroua P, Barbier F, Dieu P, et al. [Bilateral prosthesis of the knee in a hemophilia A patient with an inhibitor]. <i>Annales Francaises d Anesthesie et de Reanimation</i> 1986;5:154-6.
Lauroua P, Barbier F, Dieu P. Knee prosthesis in A haemophiliac a with inhibitor. [French]. <i>Annales Francaises d'Anesthesie et de Reanimation</i> 1986;5:154-156.

Study
Leblebisatan G, Sasmaz I, Antmen B, et al. A successful use of recombinant factor VIIa in a patient with inhibitors, for bilateral cataract operation and circumcision. <i>Haemophilia</i> 2006;12:187-189.
Leblebisatan G, Sasmaz I, Antmen B, et al. Serial urological interventions including circumcision in a hemophilic child with inhibitors. <i>Blood Coagulation and Fibrinolysis</i> 2011;22:547-548.
Lechner K, Fischer M, Santler R. Successful surgery in a patient with inhibitor substance hemophilia. [German]. <i>Wiener klinische Wochenschrift</i> 1969;81:411-416.
Leebeek FWG, Kappers-Klunne MC, Jie KSG. Effective and safe use of recombinant factor VIIa (NovoSeven) in elderly mild haemophilia A patients with high-titre antibodies against factor VIII. <i>Haemophilia</i> 2004;10:250-253.
Leggett PL, Doyle D, Smith WB, et al. Elective cardiac operation in a patient with severe hemophilia and acquired factor VIII antibodies. <i>Journal of Thoracic and Cardiovascular Surgery</i> 1984;87:556-560.
Livnat T, Barg A, Avishai E, et al. Combined bypass agents therapy for hemophilia B patients with inhibitor. <i>Haemophilia</i> 2017;23:65-66.
Livnat T, Budnik I, Levy-Mendelovich S, et al. Combination of hemostatic therapies for treatment of patients with hemophilia A and inhibitors. <i>Blood Cells, Molecules, and Diseases</i> 2017;66:1-5.
Ljung R, Petrini P, Lindgren AK, et al. Implantable central venous catheter facilitates prophylactic treatment in children with haemophilia. <i>Acta Paediatrica</i> 1992;81:918-20.
Lorenzo JI, Montoro JM, Aznar JA. Postoperative use of rFVIIa by continuous infusion in a haemophilic boy. <i>Haemophilia</i> 1999;5:135-8.
Lowe GDO, Harvie A, Forbes CD, et al. Successful treatment with prothrombin complex concentrate of postoperative bleeding in a haemophiliac with a factor VIII inhibitor. <i>British Medical Journal</i> 1976;2:1110-1111.
MacPherson BH, Drayton NE, Reid WO. Surgical hemostasis in a hemophiliac using blockade of the inhibitor system. <i>The American surgeon</i> 1974;40:224-228.
Madigan RR, Hanna WT, Wallace SL. Acute compartment syndrome in hemophilia. A case report. <i>Journal of Bone and Joint Surgery - Series A</i> 1981;63:1327-1329.
Maliekel K, Rana N, Green D. Recombinant factor VIIa in the management of a pseudotumour in acquired haemophilia. <i>Haemophilia</i> 1997;3:54-58.
Marisavljevic D, Glisic M, Elezovic I, et al. Successful extirpation of femoral pseudotumour in a patient with severe haemophilia A and an inhibitor to factor VIII. [Croatian]. <i>Srpski arhiv za celokupno lekarstvo</i> 1991;119:338-342.
Masuhara K, Shimizu T, Kawasaki N, et al. Surgery for a chronic cystic hematoma on the right thigh in a hemophiliac A with a potent factor VIII inhibitor. <i>Journal of the Japanese Orthopaedic Association</i> 1982;56:1719-1727.
Mat Husin N, Sulaiman W. Acquired factor XIII inhibitor in SLE patient: A case report. <i>International Journal of Rheumatic Diseases</i> 2016;19:57.
Matsuyama K, Ushijima K, Kano T, et al. Intraoperative use of plasma-derived activated factor VII (F VII a) in a hemophilia A patient with inhibitors. [Japanese]. <i>Japanese Journal of Anesthesiology</i> 1996;45:235-238.
McNamara JL, Lombardi JP, Ferguson R, et al. Alternative methods for anticoagulation monitoring in pediatric patients with applicability to a patient with severe hemophilia A and circulating inhibitor. <i>Journal of Extra-Corporeal Technology</i> 2001;33:239-242.
Mehta S, Nelson CL, Konkole BA, et al. Total knee arthroplasty using recombinant factor VII in hemophilia-A patients with inhibitors: A report of three cases. <i>Journal of Bone and Joint Surgery - Series A</i> 2004;86:2519-2521.
Meili EO, Dazzi H, von Felten A. [Recombinant activated Factor VII (Novoseven Novo Nordisk) for hemostasis in acquired Factor VIII-inhibitor hemophilia]. <i>Schweizerische Medizinische Wochenschrift. Journal Suisse de Medecine</i> 1995;125:405-11.

Study
Mekenkamp LJM, Beishuizen A, Slomp J, et al. Successful treatment of fulminant postoperative bleeding due to acquired haemophilia. <i>Netherlands Journal of Medicine</i> 2015;73:182-186.
Menart C, Cognet V, Petit PY, et al. Management of an anti-factor VIII inhibitor occurring during surgical procedure with continuous infusion of Novoseven. <i>Blood Coagulation & Fibrinolysis</i> 1998;9:289-90.
Menart C, Lalain JJ, Lienhart A, et al. Safety and efficacy of three arthroscopic procedures using Holmium: Yag laser in two high-responder haemophiliacs. <i>Haemophilia</i> 1999;5:278-281.
Mitchell J, Phillott A. Haemophilia and inhibitors. 2: surgical management. <i>Nursing times</i> 2008;104:28-29.
Mitrovic M, Miljic P, Darko A, et al. Sequential therapy with activated prothrombin complex concentrates and recombinant FVIIA in patients with severe haemophilia a, inhibitors and life threatening bleeding. <i>Haematologica</i> 2015;100:672.
Monteagudo J, Puig LL, Lao J. Successful operation on a haemophiliac with factor VIII inhibitor treated by plasmapheresis and factor VIII infusions. <i>Transfusion</i> 1985;25:85-86.
Montoro JB, Altisent C, Pico M, et al. Recombinant factor VIIa in continuous infusion during central line insertion in a child with factor VIII high-titre inhibitor. <i>Haemophilia</i> 1998;4:762-765.
Nabil S, Bentalha A, Jaiteh L, et al. Hydatid cyst surgery complicated by hemorrhage resistant to activated recombinant factor VII, in a hemophiliac A patient with an inhibitor. <i>Blood Coagulation and Fibrinolysis</i> 2016;27:724-726.
Nakamura M, Terashima K, Takashima Y, et al. Continuous infusion of recombinant activated factor VII during and after elbow arthroplasty in a hemophilia A patient with inhibitors. [Japanese]. [Rinsho ketsueki] <i>The Japanese journal of clinical hematology</i> 2002;43:183-188.
Nilsson IM, Berntorp E, Zettervall O. Induction of split tolerance and clinical cure in high-responding hemophiliacs with factor IX antibodies. <i>Proceedings of the National Academy of Sciences of the United States of America</i> 1986;83:9169-9173.
Nilsson IM, Jonsson S, Sundqvist SB. A procedure for removing high titer antibodies by extracorporeal protein-A-Sepharose adsorption in hemophilia: Substitution therapy and surgery in a patient with hemophilia B and antibodies. <i>Blood</i> 1981;58:38-44.
Nwachuku I, Obi N, Housden P, et al. New house or new limbs?! <i>Annals of the Royal College of Surgeons of England</i> 2010;92:W7-9.
O'Connell N, Chen J, Byrne M, et al. Massive pseudotumour resection with recombinant factor VIIa (NovoSeven) cover. <i>British Journal of Haematology</i> 2002;116:645-8.
Ogawa Y, Nishida Y, Kyo T, et al. [Successful completion of left total hip arthroplasty by inhibitor neutralization therapy in a hemophilia B patient with high responding inhibitor]. [Rinsho ketsueki] <i>The Japanese journal of clinical hematology</i> 2011;52:713-717.
O'Marcaigh AS, Schmalz BJ, Shaughnessy WJ, et al. Successful hemostasis during a major orthopedic operation by using recombinant activated factor VII in a patient with severe hemophilia A and a potent inhibitor. <i>Mayo Clinic Proceedings</i> 1994;69:641-644.
Orangio GR, Lucas GW. Management of hemophilia in colon and rectal surgery. Report of a patient with Factor VIII inhibitors and review of the literature. <i>Diseases of the Colon and Rectum</i> 1989;32:878-883.
O'Reilly RA, Hamilton RD. Acquired hemophilia, meningioma, and diphenylhydantoin therapy. <i>Journal of Neurosurgery</i> 1980;53:600-605.
Pasa S, Altintas A, Cil T, et al. Successful total hip replacement in a patient with severe haemophilia A with inhibitors using recombinant factor VIIa. <i>Haemophilia</i> 2008;14:863-5.
Pekcelen Y, Yavuz AS, Namli S, et al. Short-course use of recombinant factor VIIa in a haemophilia patient with inhibitor undergoing cataract surgeries. <i>Blood Coagulation and Fibrinolysis</i> 2005;16:445-446.

Study
Penner JA. Management of haemophilia in patients with high-titre inhibitors: Focus on the evolution of activated prothrombin complex concentrate AUTOPLEX T. <i>Haemophilia</i> 1999;5:1-9.
Perez R, Martinez RL, Pinero A, et al. Sequential treatment with bolus and continuous infusion of recombinant factor VIIa for hip arthroplasty in a patient with haemophilia A and inhibitor. <i>Haemophilia</i> 2002;8:822-5.
Pivalizza EG, Escobar MA. Thrombelastography-guided factor VIIa therapy in a surgical patient with severe hemophilia and factor VIII inhibitor. <i>Anesthesia and Analgesia</i> 2008;107:398-401.
Preston FE, Dinsdale RCW, Sutcliffe DJ. Factor VIII inhibitor by-passing activity (FEIBA) in the management of patients with factor VIII inhibitors. <i>Thrombosis Research</i> 1977;11:643-651.
Priluck IA, Howe RB, Eifrig DE, et al. Retinal surgery complicated by a spontaneously acquired factor VIII inhibitor. <i>American Journal of Ophthalmology</i> 1978;86:27-30.
Rajic N, Savic A, Popovic S, et al. Successful control of bleeding during supracondylar amputation caused by severe compartment syndrome in patient with haemophilia A and high titre of inhibitor. <i>Haemophilia</i> 2009;15:601-602.
Rana NA, Shapiro GR, Green D. Long-term follow-up of prosthetic joint replacement in hemophilia. <i>American Journal of Hematology</i> 1986;23:329-337.
Redding SW, Stiegler KE. Dental management of the classic hemophiliac with inhibitors. <i>Oral Surgery Oral Medicine and Oral Pathology</i> 1983;56:145-148.
Riaz MK, Girnius S, Palascak JE. Recombinant porcine factor VIII, OBI-1, successfully controlled gastrointestinal bleeding in a patient with acquired hemophilia A. <i>Blood</i> 2015;126 (23):4676.
Robbins D, Kulkarni R, Gera R, et al. Successful treatment of high titer inhibitors in mild hemophilia A with avoidance of factor VIII and immunosuppressive therapy. <i>American Journal of Hematology</i> 2001;68:184-188.
Rocino A, Carola A, Papa ML, et al. Major surgery for a gastric cancer in a haemophilic with high inhibitor titre successfully performed by the use of recombinant FVIIa. <i>Haemophilia</i> 1999;5:441-444.
Rodriguez-Merchan EC, Gomez-Cardero P, Jimenez-Yuste V, et al. A complex case of infected total knee arthroplasty in a haemophilic patient with inhibitor. <i>Haemophilia</i> 2012;18:e357-9.
Rutledge BP, Jasti P, Alward A. A case of acquired hemophilia preceding diagnosis of hepatocellular carcinoma. <i>American Journal of Gastroenterology</i> 2016;111:S857-S858.
Saarela MS, Tiitola M, Lappalainen K, et al. Pseudoaneurysm in association with a knee endoprosthesis operation in an inhibitor-positive haemophilia A patient - treatment with local thrombin. <i>Haemophilia</i> 2010;16:686-688.
Saba HI, Morelli GA, Azam RR, et al. Efficacy of NovoSeven during surgery on a haemophiliac with previous history of inhibitors. <i>Haemophilia</i> 2003;9:131-136.
Sadat U, Naik J, Hayes PD. Surgical complications in a hemophilia patient with factor VIII inhibitor and their endovascular management. <i>Vascular and Endovascular Surgery</i> 2008;42:168-172.
Saeki N, Mochizuki S, Fujii T, et al. Postsurgical coagulopathy in a hemophilia A patient with inhibitors: Efficacy of recombinant factor VIIa. <i>Journal of Anesthesia</i> 2014;28:621-624.
Saito R, Takahashi T, Endo H, et al. A case of subarachnoid hemorrhage complicated by acquired hemophilia. [Japanese]. <i>Neurological Surgery</i> 2009;37:1215-1219.
Salaj P, Gurlich R, Svorcova V, et al. Prophylactic preparation and surgical extirpation of a very large abdominal blood cyst in a severe haemophilia A patient with inhibitors managed by rFVIIa. <i>Haemophilia</i> 2009;15:380-382.
Santagostino E, Gringeri A, Muca-Perja M, et al. A prospective clinical trial of implantable central venous access in children with haemophilia. <i>British Journal of Haematology</i> 1998;102:1224-1228.

Study
Sartori R, Bisson R, Baars GW, et al. One-stage replacement of infected knee prosthesis in a patient with haemophilia A and high titre of inhibitors. <i>Haemophilia</i> 2008;14:375-7.
Sasaki T, Matsumoto T, Wada H, et al. Successful treatment of non-invasive bladder tumour in a haemophilia A patient with high-responding inhibitors: A case report. <i>Haemophilia</i> 2014;20:e399-e401.
Scharf R, Kucharski W, Lopaciuk S, et al. [Preliminary experience with achieving immune tolerance in patients with hemophilia A and circulating factor VIII inhibitor]. <i>Acta Haematologica Polonica</i> 1991;22:92-9.
Schulz S. Perilous complication by inhibitors of factor VIII after extraction of a tooth in a patient with haemophilia A. [German]. <i>Zeitschrift fur Klinische Medizin</i> 1985;40:533-535.
Schwartz KA, Penner JA, Fekety RF, Jr., et al. Fatal colitis in a hemophilic patient with inhibitor. <i>Southern Medical Journal</i> 1977;70:617-9.
Sessa J, Wang S, Hanna A, et al. Management of bleeding in acquired hemophilia A. <i>Critical Care Medicine</i> 2018;46 (Supplement 1):267.
Sharp HK, McIlveen LP, Schuman NJ. Use of FEIBA and Amicar in the operating room--dental treatment of a patient with hemophilia and high titer factor VIII inhibitors. Special care in dentistry : official publication of the American Association of Hospital Dentists, the Academy of Dentistry for the Handicapped, and the American Society for Geriatric Dentistry 1986;6:210-212.
Sheth S, Dimichele D, Lee M, et al. Heart transplant in a factor VIII-deficient patient with a high-titre inhibitor: Perioperative management using high-dose continuous infusion factor VIII and recombinant factor VIIa. <i>Haemophilia</i> 2001;7:227-232.
Shibata M, Nakagawa T, Akioka S, et al. Hemostatic treatment using factor VIII concentrates for neutralizing high-responding inhibitors prior to CVAD insertion for immune-tolerance induction therapy. <i>Clinical and Applied Thrombosis/Hemostasis</i> 2012;18:66-71.
Shida Y, Kondo Y, Ishikawa T, et al. A case of moderate hemophilia a with p.R1800H mutation complicated with juvenile idiopathic arthritis. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:588-589.
Shimada K, Takedani H, Inoue K, et al. Arthroscopic synovectomy of the elbow covered with rFVIIa in a haemophilia B juvenile with inhibitor. <i>Haemophilia</i> 2012;18:e414-e416.
Shobeiri SA, West EC, Kahn MJ, et al. Postpartum acquired hemophilia (Factor VIII inhibitors): A case report and review of the literature. <i>Obstetrical and Gynecological Survey</i> 2000;55:729-737.
Shurafa M, MacIntosh RB. Management of dental extractions in two hemophilia A patients with factor VIII inhibitor. <i>Journal of Oral and Maxillofacial Surgery</i> 1987;45:698-701.
Simic D, Milojevic I. The intraoperative use of recombinant FVIIa in child with hemophilia A with antibodies. <i>Paediatric Anaesthesia</i> 2007;17:789-792.
Sindet-Pedersen S, Stenbjerg S, Ingerslev J, et al. Surgical treatment of severe periodontitis in a haemophilic patient with inhibitors to factor VIII. Report of a case. <i>Journal of clinical periodontology</i> 1988;15:636-638.
Slaoui M, Lambert T, Stieltjes N, et al. Intestinal surgery with activated recombinant factor VII prophylaxis in patients with haemophilia A and high responding inhibitors: A report of five cases. <i>Blood Coagulation and Fibrinolysis</i> 2004;15:687-691.
Smirnova O, Namestnikov Y, Markova I, et al. Assessment of thrombin generation test for bypassing therapy monitoring in haemophilia patient with inhibitor. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:529.
Soker M, Zeytun H, Soker S, et al. Perioperative management and outcome of urgent appendectomy in a patient with severe hemophilia. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:847.

Study
Sorensen B, Ingerslev J. Platelet infusion supports recombinant factor VIIa in a patient with severe haemophilia A and inhibitor--clinical outcome and laboratory observations. <i>Thrombosis & Haemostasis</i> 2010;103:1275-6.
Soria A, Matichard E, Descamps V, et al. Bullous pemphigoid and acquired hemophilia. [French]. <i>Annales de Dermatologie et de Venereologie</i> 2007;134:353-356.
Stanley J, Austin SK. Use of an innovative syringe pump to deliver bolus RFVIIa for a patient with haemophilia and an inhibitor undergoing surgery. <i>Haemophilia</i> 2017;23:81-82.
Stephan B, Lauer H, Gros J, et al. Acquired hemophilia - hard to diagnose, hard to therapy, and cost-intensive! Risk-benefit ratio and critical insights based on various complications associable with a factor VIII-Dependent blood coagulation inhibitor. <i>Transfusion Medicine and Hemotherapy</i> 2015;42:31.
Stumpf UC, Eberhardt C, Kurth AA. Orthopaedic limb salvage with a mega prosthesis in a patient with haemophilia A and inhibitors - A case report. <i>Haemophilia</i> 2007;13:435-439.
Sudheesh KM, Bharani KS, Kiran HY, et al. "Antihemophilic factor is not the only answer for all factor VIII deficiencies." Case report of odontogenic infection in a patient with hemophilia A, complicated by factor VIII inhibitors, and managed by transfusion of antihemophilic factor and factor VIII inhibitor bypass activity. <i>Indian Journal of Dentistry</i> 2016;7:149-152.
Sugiura R, Kuwatani M, Kawakubo K, et al. Successful endoscopic sphincterotomy for choledocholithiasis in a patient with severe hemophilia A and inhibitors. <i>Clinical Journal of Gastroenterology</i> 2018;02:02.
Sun TZ, Lu HS, Guan ZP. Total knee arthroplasty and perioperative management of hemophilic arthritis. [Chinese]. <i>Zhonghua wai ke za zhi [Chinese journal of surgery]</i> 2007;45:708-711.
Szczepanik AB, Dabrowski WP, Szczepanik AM, et al. Sclerotherapy Of Esophageal Varices In Severe Hemophilia A Patient And High Titer Inhibitor--Case Report. <i>Polski przeglad chirurgiczny</i> 2015;87:464-468.
Tagariello G, Bisson R, Radossi P, et al. Concurrent total hip and knee replacements in a patient with haemophilia with inhibitors using recombinant factor VIIa by continuous infusion. <i>Haemophilia</i> 2003;9:738-740.
Tagariello G, De Biasi E, Gajo GB, et al. Recombinant FVIIa (NovoSeven) continuous infusion and total hip replacement in patients with haemophilia and high titre of inhibitors to FVIII: Experience of two cases. <i>Haemophilia</i> 2000;6:581-583.
Takedani H, Hirose J, Minamoto F, et al. Major orthopaedic surgery for a haemophilia patient with inhibitors using a new bypassing agent. <i>Haemophilia</i> 2016;22:e459-e461.
Takedani H, Mikami S, Kawasaki N, et al. Excision of pseudotumour in a patient with haemophilia A and inhibitor managed with recombinant factor VIIa. <i>Haemophilia</i> 2004;10:179-182.
Tamura K, Kanazawa T, Suzuki M, et al. Successful induction of immune tolerance by continuous infusion of recombinant factor VIII in a haemophilia A patient with high-inhibitor titres. <i>Haemophilia</i> 2006;12:100-102.
Tarrant J, Hughes C, Malins L, et al. Treatment of aquired factor V inhibitors resulting in disseminated intravascular coagulation. <i>Journal of Thrombosis and Haemostasis</i> 2015;13:756-757.
Thiagarajan RR, Roth SJ, Margossian S, et al. Extracorporeal membrane oxygenation as a bridge to cardiac transplantation in a patient with cardiomyopathy and hemophilia A. <i>Intensive Care Medicine</i> 2003;29:985-988.
Tourbaf KD, Dunlap BE, Ambrus JL, et al. The use of FEIBA (factor eight inhibitor bypassing activity) in cataract extraction in hemophilia A patient with inhibitor. <i>Journal of Medicine</i> 1982;13:399-410.
Tran DQ, Moss E, Murphy DA, et al. Mitral valve repair in a Jehovah's witness with haemophilia A with high-titre inhibitor. <i>Haemophilia</i> 2015;21:e523-e525.

Study
Uluca U, Soker M, Yel S, et al. Evaluation of three patients with haemorrhagic diathesis which are intracranial haemorrhage. <i>Haemophilia</i> 2015;21:84.
Vajta L, Nagy A, Kalovics J, et al. [Dental and oral surgical treatment of a B haemophilic patient with high inhibitor level. Case report]. <i>Fogorvosi Szemle</i> 2015;108:61-4.
Valentino LA, Allen G, Gill JC, et al. Case studies in the management of refractory bleeding in patients with haemophilia A and inhibitors. <i>Haemophilia</i> 2013;19:e151-e166.
Vivin P, Helmer J, Kipper J. Surgery in a patient with acquired factor VIIIC inhibitor. [French]. <i>Annales Francaises d'Anesthesie et de Reanimation</i> 1986;5:322-325.
Walsh JD, Landercasper J, Bottner WA, et al. Cholecystectomy and acquired factor VIII inhibitor coagulopathy. <i>The American surgeon</i> 2004;70:815-817.
Watts RG. Successful use of recombinant factor VIIa for emergency fasciotomy in a patient with hemophilia A and high-titer inhibitor unresponsive to factor VIII inhibitor bypassing activity. <i>American Journal of Hematology</i> 2005;79:58-60.
White B, Cotter M, Byrne M, et al. High responding factor VIII inhibitors in mild haemophilia - Is there a link with recent changes in clinical practice? <i>Haemophilia</i> 2000;6:113-115.
White IGC, Taylor RE, Blatt PM, et al. Treatment of a high titer anti-factor-VIII antibody by continuous factor VIII administration: Report of a case. <i>Blood</i> 1983;62:141-145.
Yankov IV, Spasova MI, Andonov VN, et al. Endoscopic diagnosis of intramural hematoma in the colon sigmoideum in a child with high titer inhibitory hemophilia A. <i>Folia medica</i> 2014;56:126-128.
Yildiz A, Sahin O, Yayar O, et al. An unusual coexistence of high titer inhibitor development and gastrointestinal stromal tumor in a patient with severe hemophilia: Case report. <i>Haemophilia</i> 2018;24 (Supplement 1):113-114.
Zanon E, Brandolin B, Saggiorato G, et al. Complex dental extractions in a patient with severe haemophilia A and inhibitors treated with activated prothrombin complex concentrate. <i>Blood Transfusion</i> 2012;10:225-227.
Zech R, Strother SV. Maintenance of hemostasis during exodontia in two hemophiliacs with factor VIII inhibitors. <i>Journal of Oral & Maxillofacial Surgery</i> 1983;41:53-6.

1 **Supplementary Table 7. Quality Assessment of Interventional Studies Reporting Perioperative Laboratory Monitoring**

Study	Did the trial address a clearly focused issue?	Was the assignment of patients to treatments randomised?	Were all of the patients who entered the trial properly accounted for at its conclusion?	Were patients, health workers and study personnel 'blind' to treatment?	Were the groups similar at the start of the trial?	Aside from the experimental intervention, were the groups treated equally?	How large was the treatment effect?	How precise was the estimate of the treatment effect?	Can the results be applied in your context? (Or the local population?)	Were all clinically important outcomes considered?
Furukawa S et al. 2015[20]	Y	N	Y	N	N	N	-	-	Y	N
Gatti L et al. 1984[13]	Y	N	Y	N	-	-	-	-	N	Y
Ludlam A et al. 2003[10]	Y	N	Y	N	-	-	-	-	Y	Y
Mancuso ME et al. 2016[25]	Y	N	Y	N	N	N	-	-	Y	N
Pruthi RK et al. 2007[19]	Y	Y	Y	N	N	N	-	-	Y	Y
Santagostino E et al. 2001[97]	Y	N	Y	N	-	-	-	-	Unclear	Y
Scharrer I. 1999[15]	Y	N	Y	N	-	-	-	-	N	N
Shapiro AD et al. 1998[73]	Y	Y	Y	Y	Y	Y	-	-	Y	Y

Study	Did the trial address a clearly focused issue?	Was the assignment of patients to treatments randomised?	Were all of the patients who entered the trial properly accounted for at its conclusion?	Were patients, health workers and study personnel 'blind' to treatment?	Were the groups similar at the start of the trial?	Aside from the experimental intervention, were the groups treated equally?	How large was the treatment effect?	How precise was the estimate of the treatment effect?	Can the results be applied in your context? (Or the local population?)	Were all clinically important outcomes considered?
Smith MP et al. 2001[18]	Y	N	Y	N	-	-	-	-	Y	Y

1 Y: Yes; N: No

2

1 **Supplementary Table 8. Quality Assessment of Observational Studies Reporting Perioperative Laboratory Monitoring**

Study	Was the cohort recruited in an acceptable way?	Was the exposure accurately measured to minimise bias?	Was the outcome accurately measured to minimise bias?	Have the authors identified all important confounding factors?	Have the authors taken account of the confounding factors in the design and/or analysis?	Was the follow-up of patients complete?	How precise (for example, in terms of CI and p values) are the results?
Balkan C et al. 2010[12]	Y	–	Y	–	–	Y	–
Danielson H et al. 2017[24]	Y	–	Unclear	–	–	Y	–
Habermann B et al. 2004[23]	Unclear	–	N	–	–	Y	–
Holmstrom M et al. 2012[22]	Unclear	–	Y	–	–	Y	–
Ingerslev J et al. 1996[49]	Unclear	–	N	–	–	Y	–
Kraut EH et al. 2007[14]	N	–	N	–	–	Y	–
Lauroua P et al. 2009[54]	Y	–	Y	–	–	Y	–
Mauser-Bunschoten EP et al. 1998[16]	Unclear	–	N	–	–	Y	–
Mauser-Bunschoten EP et al. 2002[17]	Y	–	N	–	–	Y	–
Negrier C et al. 2013[93]	Y	–	Y	–	–	Y	–

Study	Was the cohort recruited in an acceptable way?	Was the exposure accurately measured to minimise bias?	Was the outcome accurately measured to minimise bias?	Have the authors identified all important confounding factors?	Have the authors taken account of the confounding factors in the design and/or analysis?	Was the follow-up of patients complete?	How precise (for example, in terms of CI and p values) are the results?
Quintana-Molina M et al. 2004[65]	Y	-	Y	-	-	Y	-
Serban M et al. 2014[21]	Y	-	Y	-	-	Y	Unclear
Smith OP et al. 2002[75]	Y	-	Unclear	-	-	Unclear	-
Tjonnfjord GE. 2004, 2006 [100, 101]	Y	-	Unclear	-	-	Y	-

1 Y: Yes; N: No

2

We are submitting the full manuscript as requested.